



BEST MANAGEMENT PRACTICES FOR SOUTH DAKOTA GOLF COURSES



Table of Contents

| | |
|--|----|
| Table of Contents | i |
| Tables | v |
| Photos | v |
| Who We Are/Acknowledgements | i |
| Additional Acknowledgements | ii |
| 1 INTRODUCTION | 1 |
| 1.1 Best Management Practices | 1 |
| 1.2 Pollution Prevention | 1 |
| 1.3 Water Conservation | 2 |
| 1.4 Pollinators | 2 |
| 1.5 Creating a Facility BMP | 2 |
| 1.6 Conclusion | 3 |
| 2 PLANNING, DESIGN, AND CONSTRUCTION | 4 |
| 2.1 Regulatory Considerations | 4 |
| 2.2 Overview | 5 |
| 2.3 Wetlands and Floodplains | 6 |
| 2.4 Drainage | 10 |
| 2.5 Habitat Considerations | 11 |
| 2.6 Turfgrass Establishment | 12 |
| 2.7 External Certification Programs | 15 |
| 2.8 Planning, Design, and Construction Best Management Practices | 15 |
| 3 IRRIGATION | 18 |
| 3.1 Regulatory Considerations | 18 |
| 3.2 Water Conservation and Efficient Use Planning | 18 |
| 3.3 Irrigation Water Supply | 19 |
| 3.4 Irrigation System Design | 20 |
| 3.5 Irrigation System Installation | 22 |
| 3.6 Irrigation System Maintenance and Performance | 22 |
| 3.7 Irrigation Management Decisions | 22 |
| 3.8 Irrigation Best Management Practices | 25 |
| 4 WATER MANAGEMENT | 29 |

| | | |
|----------|--|-----------|
| 4.1 | Regulatory Considerations | 29 |
| 4.2 | Stormwater Management | 30 |
| 4.3 | Wetlands..... | 33 |
| 4.4 | Floodplains..... | 34 |
| 4.5 | Lakes and Ponds..... | 34 |
| 4.6 | Groundwater Management..... | 38 |
| 4.7 | Water Management Best Management Practices..... | 38 |
| 5 | WATER QUALITY MONITORING..... | 41 |
| 5.1 | Regulatory Considerations | 41 |
| 5.1 | Existing Water Quality Information..... | 42 |
| 5.2 | Developing a Water Quality Monitoring Program..... | 42 |
| 5.3 | Interpreting Water Quality Testing Results | 43 |
| 5.4 | Water Quality Monitoring Best Management Practices | 43 |
| 6 | NUTRIENT MANAGEMENT..... | 45 |
| 6.1 | Fertilizers Used in Golf Course Management..... | 45 |
| 6.2 | Soil pH..... | 49 |
| 6.3 | Soil Testing | 50 |
| 6.4 | Nutrient Management Planning | 50 |
| 6.5 | Fertilizer Applications..... | 54 |
| 6.6 | Nutrient Management Best Management Practices | 54 |
| 7 | CULTURAL PRACTICES..... | 56 |
| 7.1 | Mowing | 56 |
| 7.2 | Aeration..... | 58 |
| 7.3 | Surface Cultivation..... | 61 |
| 7.4 | Topdressing | 62 |
| 7.5 | Rolling..... | 63 |
| 7.6 | Cultural Practices Best Management Practices..... | 63 |
| 8 | INTEGRATED PEST MANAGEMENT | 65 |
| 8.1 | Regulatory Considerations | 65 |
| 8.2 | IPM Overview | 65 |
| 8.3 | Monitoring Pests and Recording Information..... | 66 |
| 8.4 | Identifying and Understanding Pests..... | 66 |
| 8.5 | Determining Threshold Levels..... | 70 |
| 8.6 | Control Methods..... | 70 |

| | | |
|-----------|--|------------|
| 8.7 | Evaluation and Record Keeping..... | 73 |
| 8.8 | IPM Best Management Practices | 73 |
| 9 | PESTICIDE MANAGEMENT | 76 |
| 9.1 | Regulatory Considerations | 76 |
| 9.2 | Human Health Risks..... | 77 |
| 9.3 | Personal Protective Equipment | 78 |
| 9.4 | Environmental Fate and Transport..... | 78 |
| 9.5 | Pesticide Record Keeping | 80 |
| 9.6 | Pesticide Storage and Handling | 80 |
| 9.7 | Pesticide Application Equipment and Calibration | 83 |
| 9.8 | Transportation | 84 |
| 9.9 | Mixing/Washing Station | 84 |
| 9.10 | Disposal..... | 84 |
| 9.11 | Pesticide Container Management..... | 85 |
| 9.12 | Emergency Preparedness and Spill Response..... | 85 |
| 9.13 | Pesticide Management Best Management Practices | 86 |
| 10 | POLLINATOR PROTECTION..... | 90 |
| 10.1 | Regulatory Considerations | 90 |
| 10.2 | Pest Management Practices | 91 |
| 10.3 | Preserving and Enhancing Habitat on the Course..... | 91 |
| 10.4 | Managed Bee Hives on the Course | 93 |
| 10.5 | Pollinator Protection Best Management Practices | 95 |
| 11 | MAINTENANCE OPERATIONS | 96 |
| 11.1 | Regulatory Considerations | 96 |
| 11.2 | Storage and Handling of Fertilizers | 96 |
| 11.3 | Equipment Washing | 97 |
| 11.4 | Equipment Storage and Maintenance..... | 98 |
| 11.5 | Fueling Facilities | 98 |
| 11.6 | Waste Handling | 99 |
| 11.7 | Maintenance Operations Best Management Practices | 100 |
| 12 | LANDSCAPE | 103 |
| 12.1 | Planning and Design..... | 103 |
| 12.2 | Site Inventory and Assessment | 104 |
| 12.3 | Plant Selection..... | 105 |

| | | |
|-----------|---|------------|
| 12.4 | Installation | 106 |
| 12.5 | Irrigation..... | 107 |
| 12.6 | Use of Mulch | 107 |
| 12.7 | Pruning | 108 |
| 12.8 | Pest Management | 108 |
| 12.9 | Landscape Best Management Practices | 109 |
| 13 | ENERGY | 111 |
| 13.1 | Energy Audits and Evaluation..... | 111 |
| 13.2 | Energy Efficiency Improvements..... | 112 |
| 13.3 | Green and Alternative Energy..... | 112 |
| 13.4 | Energy Best Management Practices..... | 113 |
| | REFERENCES | 114 |
| | BIBLIOGRAPHY | 118 |

Tables

| | |
|--|----|
| Table 1. Best practices for golf course planning, design, and construction..... | 7 |
| Table 2. Available soil moisture and infiltration rates for common soil textures.: | 23 |
| Table 3. Secondary macronutrients..... | 49 |
| Table 4. Soil test results and recommendations..... | 51 |
| Table 5. Annual nitrogen application rates | 53 |
| Table 6. Mowing HOC recommendations by species and location..... | 57 |
| Table 7. Aeration practices | 58 |
| Table 8. Core size options for aeration. | 59 |
| Table 9. Surface cultivation practices | 61 |
| Table 10. Light and frequent topdressing rates..... | 63 |

Photos

| | |
|--|----|
| Figure 1. Erosion control during construction is a regulatory requirement..... | 5 |
| Figure 2. Construction at Spearfish Canyon Golf Club..... | 9 |
| Figure 3. Construction at Elkhorn Ridge Golf Club..... | 9 |
| Figure 4. Bunker drainage..... | 10 |
| Figure 5. Drainage installation at Spearfish Canyon Golf Club. | 11 |
| Figure 6. Drainage installation on the 13th green at Elkhorn Ridge Golf Club. | 11 |
| Figure 7. Preparation for sodding at Sutton Bay Golf Course in Agar..... | 13 |
| Figure 8. Sod installation at Spearfish Canyon Golf Club..... | 13 |
| Figure 9. Grow-in with erosion control Sutton Bay Golf Course in Agar..... | 14 |
| Figure 10. Sod buffer around a green and bunker help to control erosion during grow-in. | 14 |
| Figure 11. Hydromulch cover on a sloped site reduces the potential for erosion..... | 15 |
| Figure 12. Advanced irrigation control systems provide precision control and monitoring. | 21 |
| Figure 13. Retention ponds temporarily hold water from storm events. | 31 |
| Figure 14. Buffer strips, such as this one at Elkhorn Ridge Golf Club, are one of the most effective means of protecting surface water quality. | 32 |

Figure 15. Wetland and pond at Spearfish Canyon Golf Club; water quality is protected by a buffer strip around the perimeter. 34

Figure 16. Algae growth in an irrigation pond. 36

Figure 17. Aeration improves dissolved oxygen levels in ponds. 37

Figure 18. Core aeration holes. 59

Figure 19. Fairway aeration. 60

Figure 20. Vertical mowing. 61

Figure 21. Topdressing. 62

Figure 22. Dollar spot. 67

Figure 23. Snow mold. 68

Figure 24. Summer patch. 68

Figure 25. Damage to turf caused by animals digging for grubs. 69

Figure 26. Controlled burning at Elkhorn Ridge Golf Club. 71

Figure 27. Controlled burns are used to manage native grass areas. 71

Figure 28. Chemical storage area at Hillsview Golf Club. 82

Figure 29. Pesticide mixing and loading using the appropriate PPE. 83

Figure 30. Milkweed plants provide habitat for monarchs. 92

Figure 31. Monarch caterpillar. 92

Figure 32. Golf courses can host managed bee hives and generate honey. 94

Figure 33. Fertilizer storage at Hillsview Golf Club. 97

Figure 34. Separating clippings reduces nutrients in washwater. 98

Figure 35. Fueling station with an AST for fuel storage. 99

Figure 36. A natural rock wall used to frame a landscaping bed adds visual interest. 103

Figure 37. Coneflowers are a favorite plant of butterflies, bees and other pollinators as well as birds. 106

Figure 38. Tree pruning using a safety harness. 108

Figure 39. Pine beetle damage. 109

Figure 40. Golf courses can become small-scale generators of energy. 112

Who We Are/Acknowledgements

Golf Course Superintendents Association of America

The Golf Course Superintendents Association of America (GCSAA) is the professional association for the men and women who manage and maintain the game's most valuable resource — the golf course. Today, GCSAA and its members are recognized by the golf industry as one of the key contributors in elevating the game and business to its current state.

Since 1926, GCSAA has been the top professional association for the men and women who manage golf courses in the United States and worldwide. From its headquarters in Lawrence, Kansas, the association provides education, information and representation to more than 17,000 members in more than 72 countries. GCSAA's mission is to serve its members, advance their profession and enhance the enjoyment, growth and vitality of the game of golf.

Environmental Institute for Golf

The Environmental Institute for Golf (EIFG) fosters sustainability by providing funding for research grants, education programs, scholarships and awareness of golf's environmental efforts. Founded in 1955 as the GCSAA Scholarship & Research Fund for the Golf Course Superintendents Association of America, the EIFG serves as the association's philanthropic organization. The EIFG relies on the support of many individuals and organizations to fund programs to advance stewardship on golf courses in the areas of research, scholarships, education, and advocacy. The results from these activities, conducted by GCSAA, are used to position golf courses as properly managed landscapes that contribute to the greater good of their communities. Supporters of the EIFG know they are fostering programs and initiatives that will benefit the game and its environment for years to come.

United States Golf Association

The United States Golf Association (USGA) provides governance for the game of golf, conducts the U.S. Open, U.S. Women's Open and U.S. Senior Open as well as 10 national amateur championships, two state team championships and international matches, and celebrates the history of the game of golf. The USGA establishes equipment standards, administers the Rules of Golf and Rules of Amateur Status, maintains the USGA Handicap System and Course Rating System, and is one of the world's foremost authorities on research, development and support of sustainable golf course management practices.

Acknowledgments

The GCSAA and EIFG wish to thank the University of Florida, Institute of Food and Agricultural Sciences, faculty, Dr. J. Bryan Unruh, Dr. Travis Shaddox, Dr. Jason Kruse, and Mr. Don Rainey, who worked on this project, providing their knowledge and expertise to help the golf course industry; the USGA for their grant to fund this important project; the volunteers who served on the task group to review BMP and provide technical assistance; and the Florida Department of Environmental Protection for permission to copy its publication, *Best Management Practices for the Enhancement of Environmental Quality on Florida Golf Courses*.



Additional Acknowledgements

The development of the *Best Management Practices for South Dakota Golf Courses* was made possible by superintendents in the state of South Dakota, turf scientists, and the South Dakota Golf Course Superintendents Association. Representatives from each organization provided their time and expertise to develop and review drafts of best management practices specifically for the state of South Dakota designed to protect the state's natural resources.

We would also like to thank the external reviewers of the draft final version of this document. The time and effort taken by individuals and agency representatives strengthened our document and ensured its accuracy.



Funding and support for this project was provided by the Golf Course Superintendents Association of America (GCSAA), the Environmental Institute for Golf (EIFG) and the United States Golf Association (USGA). The GCSAA also provided a Best Management Practices template that served as source material for this project.

1 INTRODUCTION

South Dakota's golf course superintendents are dedicated to protecting the state's natural resources. As a demonstration of this commitment, superintendents have partnered with turf scientists to develop and document best management practices (BMPs) for golf course management. These research-based, voluntary guidelines have been developed specifically for South Dakota's golf courses. These guidelines not only protect natural resources, they also afford the opportunity for superintendents to be recognized as environmental stewards by club members, the community at large, and state officials.

Golf courses in the state's urban areas, for example, allow water to infiltrate into the ground naturally instead of flowing into storm sewers or streams and rivers. Because of the concerns related to nutrient enrichment of the Missouri River watershed, being proactive stewards of water quality resources in this and any watershed is especially important. Golf courses provide additional environmental benefits to the public, such as creating wildlife habitat.

1.1 Best Management Practices

BMPs are methods or techniques found to be the most effective and practical means of achieving an objective, such as preventing potential negative water quality impacts or reducing pesticide usage. Because of the efforts aimed at protecting surface and groundwater quality, the majority of BMPs in this document relate to water.

Many BMPs protect water quality by reducing nonpoint source pollution (such as nutrients and pesticides in stormwater runoff), stormwater volume, and peak flow. Through retention, infiltration, filtering, and biological and chemical actions, preventing or minimizing the effects of golf course management on surface and groundwater resources is easily achievable. In fact, several studies have shown that implementing BMPs improves water quality as it traverses golf course properties. Many BMPs also can be used to conserve our water resources and to prepare for water use restrictions that may be imposed in times of extended drought.

1.2 Pollution Prevention

Potential contaminants in water have one of two sources: point source or nonpoint source. A point source is a discharge from a discrete location (such as a pipe); nonpoint source pollution is caused by water moving over and through the ground picking up and carrying away natural and human-made pollutants and finally depositing them into lakes, rivers, wetlands, and ground waters.

Best management practices reduce the potential for nonpoint source impacts to water quality include sedimentation, runoff, leaching, and drift. For example, bare ground is much more likely to erode than turf. Therefore, following grow-in BMPs during course construction or renovation to quickly establish dense turf ground cover helps minimize erosion potential. Maintaining vegetated areas, such as filter strips and buffers, which slow down stormwater and any excess

irrigation, allows soil infiltration and plant uptake and is another key BMP. Pesticide BMPs help superintendents follow state and federal regulations related to the storage, handling, transport, and use of pesticides, preventing spills from becoming point sources of pollution and minimizing the potential for nonpoint source pollution from these chemicals.

Understanding golf course site characteristics is another key to preventing pollution. Some sites, such as the Missouri River Valley slopes in eastern South Dakota, are more prone to runoff. The sandhills of southwestern South Dakota are more prone to leaching due to sandy soils and shallower water tables.

1.3 Water Conservation

Water is a fundamental element for physiological processes in turfgrasses such as photosynthesis, transpirational-cooling, as well as for the diffusion and transport of nutrients. Turfgrass quality and performance depend on an adequate supply of water through either precipitation or supplemental irrigation. Too little water induces drought stress and weakens the plant, while excess soil water causes anaerobic soil conditions that stunt plant growth and promote disease. Excessive water can also lead to runoff or leaching of nutrients and pesticides into groundwater and surface water. Proper irrigation scheduling, careful selection of turfgrass species, and incorporation of cultural practices that increase the water holding capacity of soil are addressed through these BMPs, as well as considerations in the design, construction, operation and maintenance of irrigation systems.

1.4 Pollinators

Protecting bees and other pollinators is important to the sustainability of agriculture. Minimizing the impact of pesticides on bees, other pollinators, and beneficial arthropods is addressed in this document in two ways: providing specific guidance for pesticide applicators and promoting the use of integrated pest management (IPM) methods to reduce pesticide usage and minimize the potential of exposure. Superintendents can also directly support healthy pollinator populations by providing and/or enhancing habitat for pollinator species and by supplying food sources and nesting sites and materials.

1.5 Creating a Facility BMP

To adapt BMPs to an individual facility, superintendents should assess their individual site, consider their available resources (such as budget and staffing), and understand that implementing BMPs will be an ongoing process. In addition, understand that implementing BMPs will be a process that can be undertaken over time and that multiple approaches can successfully protect natural resources. Golf course quality and playability does not suffer while implementing BMPs. BMPs focus on golf courses being managed in harmony with the natural environment.

Besides contributing to natural resources stewardship, incentives for golf courses in South Dakota to create a facility BMP plan and to implement BMPs include the following:

- Potential for more efficiently allocating resources by identifying management zones.
- Cost savings associated with changing from a potential overreliance upon fertilizers and pesticides to applying less of each.
- Cost savings associated with more efficient irrigation and other water conservation efforts.
- Improved community relations.
- Recognition by club members and the community at large as environmental stewards.

Because of limitations, such as budget, staff, clientele expectations, and management decisions, not all golf courses can achieve all of the best practices. It should also be recognized that golf courses are often managed with BMPs at the forefront of the superintendent's annual program. Yet documenting those efforts in a cohesive format has often been lacking. This BMP program for South Dakota golf courses will serve as a resource and a catalyst for tracking compliance with up to date research-based BMPs. However, planning for improvements over time and making small changes that meet the goals of BMPs can be achieved. For example, while a sophisticated washwater recycling system may be too expensive for many facilities, blowing clippings off mowers onto a grass surface is easily achieved and markedly reduces the amount of nitrogen and phosphorus in clippings that end up in washwater.

1.6 Conclusion

This document was developed using the latest science-based information and sources. This resource is intended to be a living document. Therefore, the latest version of this document will be posted on the South Dakota GCSAA website (<https://www.sdgcga.org/>). The South Dakota BMP steering committee intends to review this information periodically. As of the time of this publication, the information was the most recent available. Some sources are updated regularly, and the reader should try to identify the latest version. In addition, regulations may change, and the reader should identify any changes since the publication date.

2 PLANNING, DESIGN, AND CONSTRUCTION

Building a new golf course or renovating an existing one requires careful protection of natural resources during all phases of planning, design, and construction. Implementing BMPs should help achieve efficiencies in operation and management as well as an environmentally sustainable golf course.

2.1 Regulatory Considerations

Regulations are in place at the local, state, and national levels that impact planning, design, and construction activities on South Dakota's golf courses. These laws are in place to protect and conserve the environment both during and after the construction process. Before beginning any golf course construction or renovation, consultation with the appropriate regulatory agencies is recommended. Communication between developers, designers, owners, the public, and regulatory agencies should occur early and often. In addition, if new wells must be installed, experts should be consulted for proper siting in the design plan, and all setback and other regulatory requirements.

2.1.1 Stormwater Permits

The Environmental Protection Agency (EPA) protects streams, rivers and lakes from construction pollution under the Clean Water Act (CWA). The South Dakota Department of Environment and Natural Resources (DENR) creates state-specific regulations. A [National Pollutant Discharge Elimination System \(NPDES\) permit](#) issued by DENR is required for any construction-related disturbance greater than one acre.

The development of a Stormwater Pollution Prevention Plan (SWPPP) that addresses erosion and sediment control during construction is a primary condition of stormwater construction NPDES permits. The SWPPP must be kept on site and available for inspection. More information on SWPPPs can be found on the EPA's [Developing a Stormwater Pollution Prevention Plan \(SWPPP\)](#) webpage.

2.1.2 Wetlands

Activities that impact wetlands are regulated under sections 404 and 401 of the federal CWA. The U.S. Army Corps of Engineers (USACE) regulates dredging and filling of waters in the United States under Section 404 of the CWA. The DENR administers the [Section 401 Water Quality Certification Program](#) of the CWA. Consultation with DENR during the design phase of any construction activities expected to impact wetlands will assist in understanding the permitting process.



Figure 1. Erosion control during construction is a regulatory requirement.

2.1.3 Water Usage

DENR is the state agency authorized to regulate surface and groundwater usage. All diversions of surface and groundwater for irrigation, hydropower, industrial use, municipal use, storage, and other uses require a [state permit](#). Reporting requirements are identified in permits. Superintendents should ensure that required reports are submitted on time as required.

2.1.4 Listed Species

A number of federally listed threatened and endangered species can be found in South Dakota, along with candidate species. The South Dakota Ecological Services program, part of the United State Fish and Wildlife Service, provides information on federally listed and candidate species and their critical habitat and provides [species list by county](#). Information on rare state plant and animal species is maintained by the [South Dakota Natural Heritage Program](#). South Dakota Game, Fish and Parks can conduct [environmental reviews](#) prior to construction.

2.2 Overview

Proper planning is the first step to any construction or renovation project, minimizing expenses from unforeseen events and maximizing long-term success. Good planning also incorporates

conservation of natural resources into the project. The design should allow for economic sustainability while meeting the stakeholder needs. Once a course is designed, construction must be carried out in a way that minimizes environmental impacts. Maintaining a construction progress report, as well as following regulations and coordinating with regulatory agencies as required, helps to ensure compliance. Table 1 summarizes the steps and best practices for each phase of the planning, design, and construction process. Topics of particular importance, such as wildlife habitat, erosion and sedimentation control, drainage, and grow-in, are then discussed in more detail.

2.3 Wetlands and Floodplains

2.3.1 Wetlands

The State of South Dakota has adopted the following federal definition of wetlands:

“Land that has a predominance of hydric soils; is inundated or saturated by surface or ground water at a frequency and duration sufficient to support a prevalence of hydrophytic vegetation typically adapted for life in saturated soil conditions; and under normal circumstances does support a prevalence of such vegetation.”

Most of South Dakota’s wetlands are largely referred to as “prairie potholes” -- natural depressions in the landscape, courtesy of retreating glaciers 12,000 years ago. Potholes vary in size from a fraction of an acre to hundreds of acres. In South Dakota, depressional wetlands are found primarily east and north of the Missouri River. For more information on these eastern wetlands, see the publication [Eastern South Dakota Wetlands](#). Wetlands occur in the western portion of the state but are less numerous. Conserving the state’s wetlands protects water quality and biodiversity, while reducing the potential for flooding and soil erosion.

To protect this natural resource, wetlands should be identified in the field by qualified wetland specialists during the design phase and before the permitting process is initiated. Course design should minimize any impact to wetlands and streams tied to activities such as filling, dredging, flooding, crossings, or converting areas from one habitat type to another. In addition, natural buffers should be retained around wetlands (as with other waterbodies) to protect water quality and provide habitat.

In some instances, wetlands and streams can be improved or restored during golf course construction. For example, a highly degraded stream or wetland can sometimes be reshaped, rehabilitated, or replaced entirely to meet project goals and improve ecological function. Qualified environmental consultants can evaluate the overall benefit of stream enhancement or restoration and assist with permitting issues, which may include a federal 404 permit and/or state 401 certification.

Table 1. Best practices for golf course planning, design, and construction

| Planning | |
|--|--|
| Step | Description |
| <i>Assemble Team</i> | The team should include, but not be limited to, a golf course architect, golf course superintendent, clubhouse architect, irrigation engineer, environmental engineer, energy analyst, economic consultant, civil engineer, soil scientist, golf course builder, and a legal team. |
| <i>Define Objectives</i> | Identify realistic goals, formulate a timeline, etc. |
| <i>Conduct a Feasibility Study</i> | Evaluate finances, environmental issues, water availability and sources, and energy, materials, and labor needs. Identify applicable government regulations. |
| <i>Select Site</i> | Site should meet project goals and expectations. Identify all strengths and weakness of each potential site. During site selection, any site constraints, such as the presence of listed species or valuable habitat, should be identified. |
| Design | |
| <i>Retain a Project Manager/Superintendent</i> | This person is responsible for integrating sustainable practices in the development, maintenance, and operation of the course. |
| <i>Design the Course</i> | Existing native landscapes should remain intact as much as possible. Should consider adding supplemental native vegetation to enhance existing vegetation alongside lengthy fairways and out-of-play areas. Nuisance, invasive, and exotic plants should be removed and replaced with native species adapted to the area. |
| | Greens: Should have plenty of sunlight and be well drained. Greens should be big enough to have several multiple hole cupping locations that can handle expected traffic. |
| | Root zone material should be selected with United States Golf Association (USGA) specifications in mind, as published in A Guide to Constructing The USGA Putting Green . |
| | Grass Selection: Species should be selected based on climate, environmental, and site conditions and species adaptability to those conditions, including disease resistance, drought tolerance, spring greenup, and traffic tolerance. |
| | <p>Bunkers: The number and size of bunkers depends on considerations, such as the resources available for daily maintenance. For each bunker consider:</p> <ul style="list-style-type: none"> • The need for drainage. • Entry/exit points and how these will affect wear-and-tear patterns. • The proper color, size, and shape of bunker sands to meet needs. <p>New bunker construction techniques can also be researched to see if they satisfy stakeholders' needs.</p> |

| | |
|-------------------------------------|---|
| <i>Design Irrigation System</i> | Hire a professional irrigation architect, if possible, to design the irrigation system. Keep in mind the different water needs of greens, tees, fairways, roughs, and native areas. Consider the topography, prevalent wind speeds, and wind direction when spacing the heads. Choose the most efficient type of irrigation system considering available resources. The “Irrigation” chapter of this document provides detailed information on irrigation-related BMPs. |
| Construction | |
| <i>Select Qualified Contractors</i> | Use only qualified contractors who are experienced in the special requirements of golf course construction. Members of the Golf Course Builders Association of America make great candidates. |
| <i>Safeguard Environment</i> | Follow all design phase plans and environmental laws. Soil stabilization techniques should be rigorously employed to maximize sediment control and minimize soil erosion. Temporary construction compounds and pathways should be built in a manner that reduces environmental impacts. |
| <i>Install Irrigation System</i> | Installation should consider the need to move equipment and bury pipe while maintaining the original soil surface grade to minimize the potential for erosion. |
| <i>Establish Turfgrass</i> | Turfgrass establishment methods and timing should allow for the most efficient progress of work, while optimizing resources and preventing erosion from bare soils before grass is established. |



Figure 2. Construction at Spearfish Canyon Golf Club.



Figure 3. Construction at Elkhorn Ridge Golf Club.



Figure 4. Bunker drainage.

2.3.2 Floodplains

Golf course development is often compatible with floodplains, particularly when compared with other uses such as residential or commercial development. Minimizing encroachment into floodplains to the extent possible is prudent. Any substantial disturbance to a floodplain, including clearing and grading, generally requires an engineering analysis to demonstrate minimal impact on the base flood elevation in accordance with local ordinances. Depending on the complexity of the encroachment, this analysis may be as simple as a comparison of cut and fill quantities within the floodplain or as complex as a detailed floodplain model of the entire watershed. A complex analysis may require a Federal Emergency Management Agency (FEMA) review along with potential revision to the floodplain mapping.

Key course components (such as greens and tees) should be designed above the 100-year flood elevation whenever possible to avoid loss of golf play due to periodic flooding. Any effects on the floodplain and floodway should be considered, and the required offsetting adjustments should be made in grades or vegetative treatment.

2.4 Drainage

Adequate drainage is necessary for healthy turfgrass. Average rainfall and frequency guides the design of drainage systems. For example, a golf course drainage system is typically designed to retain a two- or five-year rain event, reasonably draining the precipitation in a matter of hours; excess water that has not infiltrated into the ground is retained temporarily until it leaves the property. In some instances, golf courses are mandated to handle a 20-, 50-, or 100-year rain

event; this ability to retain large amounts of water requires accurate engineering and more extensive construction.



Figure 5. Drainage installation at Spearfish Canyon Golf Club.

Many BMPs prolong the retention process as long as practical, retaining as much of the stormwater in surface or underground storage as is reasonable, and may even improve the quality of water leaving the property. A high-quality BMP plan for drainage addresses runoff containment, adequate buffer zones, and filtration techniques. However, drainage of golf course features is only as good as the system's integrity. Damaged, improperly installed, or poorly maintained drainage systems negatively impacts play and increases risks to water quality.

2.5 Habitat Considerations

In urban and suburban environments, a golf course may provide the best habitat for many species. A number of golf course management activities can maintain and enhance habitat, and provide food and shelter for numerous species,



Figure 6. Drainage installation on the 13th green at Elkhorn Ridge Golf Club.

including mammals, birds, fish, amphibians, reptiles, insects, and native plants. Examples of ways to maintain and enhance habitat include:

- Identifying and preserving wildlife and migration corridors to help maintain populations at sustainable levels.
- Retaining natural buffer areas around wetlands and watercourses to preserve habitat while protecting water quality for aquatic species.
- Planting native species to provide food for animals and insects.
- Retaining dead trees to serve as nesting areas.
- Providing nest boxes for birds, bees, and bats.
- Removing exotic and invasive species to improve habitat.

The “Pollinator Protection” and “Landscape” chapters provide additional recommendations and BMPs for enhancing habitat on the golf course. The South Dakota Department of Agriculture (SDDA) publishes a [noxious weed and pest list](#) and a [weed watch list](#) that can provide useful information for invasive species control efforts.

2.6 Turfgrass Establishment

2.6.1 Species Selection

South Dakota’s climate is generally favorable to cool season grasses (fine fescue, Kentucky bluegrass, perennial ryegrass, and tall fescue), with the exception of the warm season buffalograss, a Great Plains native. When selecting species and cultivar, site specific characteristics, such as desired use, site and microclimate conditions, disease resistance, drought tolerance, and spring transition traits, should be considered.

Buffalograss (*Buchloe dactyloides*) can be used as a low-maintenance turfgrass in South Dakota. It forms a dense sod that establishes quickly through stolons, has good color, and has exceptional drought, heat, and cold tolerance -- qualities that can benefit the golf industry by reducing management inputs and costs. These inputs, such as reduced nitrogen requirements, are significantly less than those used to manage most other turf species, such as Kentucky bluegrass. Although buffalograss requires minimal inputs once established, it takes longer to establish than cool season grasses and requires a higher level of maintenance during establishment to ensure an adequate stand. As a warm-season grass, it has a short growing season, greening up in mid to late May and staying green until the cooler temperatures and frost cause it to lose its green color and turn brown in mid to late September. In addition to proper fertilization, weed control during the first year is essential to ensure adequate establishment because high weed pressure at this time can greatly reduce quality and density for several years. Newer buffalograss cultivars are being developed that are superior to common buffalograss for turfgrass quality, uniformity, production characteristics, and canopy density.

2.6.2 Seedbed Preparation and Planting

Proper seedbed site preparation can help avoid long-term problems, such as weed encroachment, diseases, and drought susceptibility. Debris should be removed that could hinder root growth and limit access to water and nutrients. Any drainage issues should be corrected through grading and installation of drainage technologies.

Turfgrass gets off to the best start when proper attention is given to:

- Preparing the site and soil.
- Understanding correct planting techniques for the material being used (seed, sod or plugs).
- Properly caring for the grass after planting.

Industry sales representatives are also good resources for the latest in available turfgrass cultivars.



Figure 7. Preparation for sodding at Sutton Bay Golf Course in Agar.



Figure 8. Sod installation at Spearfish Canyon Golf Club.

2.6.3 Erosion and Sediment Control

The loss of topsoil from a site can be a problem for numerous reasons. Soil carried by wind and water transports contaminants with it. For example, erosion can enrich surface waters, where phosphorus, and to a lesser extent nitrogen, can cause eutrophication (the enrichment of waterbodies by inorganic plant nutrients). When sediments and soils enter water, they can also increase turbidity, which can have harmful effects on aquatic plants and animals. Therefore, control measures should be documented in an erosion and sediment control plan, put in place prior to any soil disturbance, and properly maintained.



Figure 9. Grow-in with erosion control Sutton Bay Golf Course in Agar.



Figure 10. Sod buffer around a green and bunker help to control erosion during grow-in.



Figure 11. Hydromulch cover on a sloped site reduces the potential for erosion.

2.7 External Certification Programs

Golf-centric environmental management programs or environmental management systems, such as Audubon International’s [Audubon Cooperative Sanctuary Program for Golf](#) and the Groundwater Foundation’s [Groundwater Guardian Green Sites](#) program, can help golf courses protect natural resources, as well as gain recognition for their environmental education and certification efforts.

2.8 Planning, Design, and Construction Best Management Practices

Planning and Design

- ❖ Maintain appropriate silt fencing during construction to prevent erosion and sedimentation in accordance with the SWPPP.
- ❖ Establish a low- to no-maintenance level within a 75-foot buffer along wetlands during construction.
- ❖ Establish and maintain a 100-foot riparian buffer around wetlands, springs, and channels during construction.

Wetlands and Floodplains

- ❖ Install stream buffers to restore natural water flows and flooding controls.
- ❖ Install buffers in play areas to stabilize and restore natural areas that attract wildlife.
- ❖ Install retention basins to store water and reduce flooding at peak flows.

Drainage

- ❖ Evaluate the watershed size to understand drainage needs and appropriate pipe sizing.
- ❖ Ensure that no discharges from pipes go directly to surface water.
- ❖ When constructing drainage systems, pay close attention to engineering details such as subsoil preparation and the placement of gravel, slopes, and backfilling.
- ❖ Discharge internal golf course drains through pretreatment zones and/or vegetative buffers to help remove nutrients and sediments. Do not discharge directly into an open waterbody.
- ❖ Routinely inspect the drainage system to ensure proper function.

Listed Species

- ❖ Identify any listed species and critical habitat that may be present on the site and preserve habitat, including feeding and nesting areas.

Habitat Conservation

- ❖ Identify the different types of habitat specific to the site.
- ❖ Identify habitat requirements (food, water, cover, space) for wildlife species.
- ❖ Identify and preserve regional wildlife and migration corridors by avoiding or minimizing crossings. Design unavoidable crossings to accommodate wildlife movement.
- ❖ Design and locate cart paths to minimize environmental impacts. Construct the paths with permeable materials, if possible.
- ❖ Remove nuisance and exotic/invasive plants and replace them with native species that are adapted to a particular site.
- ❖ Maintain clearance between the ground and the lowest portion of any fences or walls to allow wildlife to pass, except in areas where feral animals need to be excluded.
- ❖ Retain dead tree snags for nesting and feeding sites, provided they pose no danger to people or property.
- ❖ Construct and place birdhouses, bat houses, bee boxes, etc. in out-of-play areas.
- ❖ Plant pollinator gardens around the clubhouse and out-of-play areas.
- ❖ Retain riparian buffers along waterways to protect water quality and provide food, nesting sites, and cover for wildlife.

Turfgrass Establishment

- ❖ Select species and varieties that are adapted to the desired use, taking note of disease resistance, spring transition and greenup, drought tolerance, and other traits such as shade and wear tolerance.
- ❖ Prepare seed/sod beds to maximize success.
- ❖ Ensure erosion and sediment control devices are in place and properly maintained.
- ❖ Plant cool season grasses from seed from mid-August to early September to allow the seed to germinate and develop well before cold temperatures significantly slow growth prior to winter.
- ❖ Plant warm-season grass seed when the temperature is 70 to 95 degrees F.).

- ❖ Plant or establish sod when the turfgrass is actively growing so the sod will root or “knit” down into the soil as quickly as possible
- ❖ Fill gaps in sod seams with soil or sand to provide a uniform surface.
- ❖ Use selective pre-emergence herbicides to reduce weed competitions and improve the chance of success with seeding establishment during the spring.
- ❖ Apply a fertilizer containing phosphorus at seeding. An additional application should be applied if turf displays symptoms of phosphorus deficiency.
- ❖ Nitrogen and sufficient water are essential during establishment. Light and frequent applications of nutrients are most desirable, unless a slow-release nitrogen source is applied.
- ❖ Allow the turfgrass to initially grow one-third to one-half higher than the desired mowing height before beginning to mow and never remove more than one-third of the turf leaf at mowing.
- ❖ Reducing watering prior to mowing will help the soil dry a bit to better tolerate the weight of the mower.
- ❖ Consider mowing with a walk behind mower rather than a heavier riding mower to avoid making wheel track depressions in the soil.
- ❖ Keep mower blades sharp. Dull mowers may dislodge or damage young seedlings.

3 IRRIGATION

The supplemental use of water in turfgrass management encourages healthy turfgrass and landscape plants. BMPs related to irrigation conserve and protect water resources. Conservation and efficiency-related efforts consider the strategic use of course and irrigation design, computerized and data-integrated scheduling, and alternative water quality/supply options that maximize plant health and reduce the potential for negative impacts on natural resources.

Irrigation BMPs may also provide an economic, regulatory compliance, and environmental stewardship advantage to those who integrate them into an irrigation management plan. BMPs are not intended to increase labor or place an undue burden on the owner or superintendent. Applied appropriately, irrigation-related BMPs can help stabilize labor costs, extend equipment life, reduce repairs, and limit overall personal and public liability while protecting and conserving natural resources.

3.1 Regulatory Considerations

DENR manages the use of surface and groundwater in the state. A [permit](#) is required to withdraw water for irrigation reporting requirements. DENR provides guidance for [drilling wells](#).

During times of extended drought, groundwater use restrictions may be issued or imposed by municipal governments. Superintendents should be aware of water use restrictions and be prepared with a drought management plan.

The “Water Management” chapter of this document provides additional information on regulatory policies and agencies that regulate the use and protection of South Dakota’s water resources.

3.2 Water Conservation and Efficient Use Planning

Potable water supplies in many areas of the United States are limited, and demand continues to grow. The challenge is to find solutions to maintain the quality course conditioning while using less water. Opportunities to conserve water exist during initial course design and renovations, during irrigation system design and use, and by incorporating the use of management zones of different inputs.

Some courses are designed using a “target golf” concept that minimizes the acreage of irrigated turf. If properly designed, water hazards and stormwater ponds can capture rain and runoff that may provide supplemental irrigation water storage in ponds or lakes under normal conditions. Backup sources may be needed during severe drought. During times of intense heat stress, syringing, which is the practice of applying a small amount of water to the turfgrass leaf canopy without adding significant amounts of water to the soil, help cool the turfgrasses as it evaporates from the leaf canopy, may be beneficial under certain conditions. These conditions include turf with a very shallow root system and turf compromised by disease, poor soils, or wet-wilt.

Because the evaporative cooling effect of syringing is temporary, repeated syringing and/or the use of fans will maximize the cooling effect.

In addition to utilizing well-adapted cultivars for in-play areas, existing golf courses can convert out-of-play area turf to naturally adapted native plants, grasses, or ground covers to reduce water use and augment the site's aesthetic appeal. Native plant species also provide wildlife with habitat and food sources, such as native flower areas that benefit pollinators. After establishment, site-appropriate plants normally require little to no irrigation. See the "Pollinator Protection" and "Landscape" chapters of this document for more information on native and drought-tolerant plants.

Water conservation plans should be prepared before mandatory water restrictions are enacted in times of drought. These plans should identify ways to achieve a 10% reduction in water use. Communication with water managers, golf club members, and the public should be maintained to explain water conservation efforts as a proactive approach to addressing water-related issues.

3.3 Irrigation Water Supply

3.3.1 Irrigation Water Sources

Golf course designers and managers should identify and use alternative water supply sources to conserve freshwater drinking supplies whenever possible. The routine use of potable water is not a preferred practice; therefore, municipal drinking water should be considered only when no alternatives exist.

3.3.2 Irrigation Water Quality

Irrigation water should be assessed to determine its suitability for irrigation and for plant growth. Non-potable water irrigation sources (such as retention ponds and recycled water) should be regularly tested to ensure that the quality is within acceptable parameters for irrigation. The assessment should identify the chemical characteristics of the water and address possible issues with soil salinity and plant health caused by poor water quality.

For more information, on irrigation water quality specifically for turfgrass, see:

- [*Irrigation Water Quality Guidelines for Turfgrass Sites*](#), Penn State University.
- [*Understanding Water Quality and Guidelines to Management*](#), USGA Green Section.

3.3.3 Irrigation Water Requirements

An adequate water supply is essential for any planned or proposed golf course irrigation system. Water requirement analyses can be conducted to understand irrigation needs. A seasonal bulk water requirement analysis verifies the suitability of a water source and irrigation system to supply irrigation water under normal conditions. Because of the potential for drought, a maximum seasonal bulk water requirement analysis should be conducted as a worst-case

scenario and calculated to not account for rainfall. For more information on calculating water requirements and example calculations, see Chapter 3 of [*Environmental Best Management Practices for Virginia's Golf Courses*](#).

It is essential that all delivery systems install and maintain accurate metering devices. Being able to measure water use allows baselines to be established and progress in water conservation efforts to be tracked. Installation of water meters will become more critical as more regulatory and compliance obligations are imposed on water for irrigation.

3.4 Irrigation System Design

3.4.1 Site Assessment

An assessment of the entire facility should be conducted prior to developing a system design. The site assessment should include site-specific features, such as water sources; soil types (see the [Web Soil Survey](#) for identifying site-specific soil types), soil physical properties (Table 3-1); microclimates; slopes; sun, wind and shade exposures; and a seasonal and bulk water requirement analysis. In addition, the site assessment should evaluate the impact of design elements, such as design features and concepts, planned or existing turfgrass species and varieties, and planned or existing drainage systems. The system design should include a general irrigation schedule with recommendations and instructions on modifying the schedule to meet these site-specific needs.

3.4.2 Design

A well-designed irrigation system should operate at peak efficiency and be designed and installed to optimize water use efficiency, focusing on water placement and distribution. The design should maximize water use, reduce operational cost, conserve supply, and protect water resources. Where feasible, variable frequency drive (VFD) pumps and/or pump station should be used. These systems only expend enough energy to meet the demands of the irrigation pump(s). VFD systems reduce water hammer to fitting, pipe, and sprinklers when systems are pressurized. Detailed BMPs for irrigation system design are published by the Irrigation Association in [2014 Landscape Irrigation Best Management Practices](#).

The irrigation system design should meet the site-specific needs identified by the water quantity analyses and the site assessment. The system's capacity to deliver water should not exceed the infiltration of the soils on site, as that will lead to runoff. Though the design of an irrigation system is complex, some of the most important design decisions that influence the efficiency and effectiveness of water usage include those related to sprinkler and piping placement, sprinkler coverage and spacing, and communication options.

Multi-row sprinkler systems provide the most efficient use of water and can respond to specific moisture requirements of selected areas. Newer designs with multiple nozzle configurations provide increased flexibility and improved distribution uniformity. Single row systems do not

uniformly distribute water and increases the risk of runoff. Double-row systems offer improved efficiency over single-row coverage, although manual watering or other types of supplemental watering may be needed outside the fairway area and into the extended rough. Sprinkler layouts can be specific to each area. For example, part-circle sprinklers can be arranged to avoid overspray of impervious surfaces and to apply water only to the green surface or in heavy traffic areas. Manual quick-coupler valves can be an important conservation element and should be installed near greens, tees, and bunkers so these can be hand-watered during severe droughts. Irrigation systems designs strive to provide uniform water distribution and to achieve distribution uniformity (DU) values near 80%. After installation, nozzles and irrigation head runtimes should be optimized to maintain uniform soil moisture distribution. That can be monitored with a soil moisture probe.

For precise irrigation control, courses should consider using advanced irrigation control systems that can schedule each green, tee, and fairway separately and allow course managers to adjust for differences in microclimates and root zones. Weather stations that calculate and automatically program water replacement schedules also provide opportunities for more precise irrigation, as do soil moisture sensors placed in multiple locations. Additional features may include rain stop safety switches that either shut down the system in the event of rain or adjust schedules based on the amount of precipitation.



Figure 12. Advanced irrigation control systems provide precision control and monitoring.

3.5 Irrigation System Installation

To ensure maximum efficiency, the irrigation system must be installed per the design and specifications. The installer must ensure that there is qualified supervision of the installation process and that a qualified irrigation specialist inspects and approves the system installation.

3.6 Irrigation System Maintenance and Performance

A properly working irrigation system is critical to ensure optimum operation. System checks and routine maintenance should be done for pumps, valves, programs, fittings, and sprinklers. A schedule of inspections and a plan for record keeping should be completed. Use of photography is especially helpful in recording installations/repairs of underground systems. The publication [2014 Landscape Irrigation Best Management Practices](#) can be consulted for devising a schedule and a plan for record keeping. To ensure that it is performing as intended, an irrigation system should be calibrated regularly by conducting periodic irrigation audits, such as catch-can tests and an annual irrigation audit, to check actual water delivery and nozzle efficiency. Nozzles can wear over time. This will change irrigation output and distribution. Nozzles should be replaced, depending on the manufacturer's recommendation, to ensure proper function.

While routine inspection and audits can be performed by the golf course superintendent, a professional irrigation consultant is required for a detailed irrigation audit, which should be conducted according to the [Irrigation Audit Guidelines](#) published by the Irrigation Association. Ideally, this professional audit should be conducted at least once every five years.

3.6.1 Seasonal Maintenance

Winterizing protects irrigation system pipes from damage due to water expanding and rupturing the pipe walls and fittings. Golf courses need to drain or use compressed air to remove water from lateral and mainline pipes to prepare for when temperatures drop below freezing. Many facilities operate an independent irrigation system below the frost line, allowing the facility to apply water during cold periods to dormant turf in an effort to prevent winter desiccation and winter kill, as described in the "Irrigation Scheduling" section of this chapter (Section 3.7.5).

3.7 Irrigation Management Decisions

An irrigation system program and run times should be operated based only on the moisture needs of the turfgrass. Additionally, irrigation may be needed to satisfy "water-in" requirements for a fertilizer or chemical application as directed by the label. Calendar or seasonal considerations are only a general starting point in irrigation programming. Therefore, irrigation scheduling must also consider soil infiltration and percolation rates, as well as plant water requirements, to prevent excessive irrigation that could lead to leaching and runoff and poorer turfgrass quality.

3.7.1 Soil Infiltration Rate and Plant Available Water

The infiltration rate depends on soil types and textures. Sandy soils, with their higher porosity, have greater infiltration rates than silty or clay soils. Plant available water (PAW) represents the amount of water (expressed in inches) available per inch of soil depth that a plant can extract for transpiration, given a healthy root system through a particular depth of soil. A soil moisture probe indicates the total volumetric water content, which is greater than the PAW for a soil. The PAW can be estimated with a soil moisture meter by subtracting the current soil moisture content from the moisture content when the turf wilts. Plant available soil moisture and infiltration rates are provided in Table 2.

3.7.2 Root Zone Depth

The effective turf rooting depth should be determined with a soil probe or spade. Golf greens and tees have the majority of roots in the top several inches of soil, while fairways and roughs will typically have deeper roots. Exact root depths depend on grass species, the soil environment (soil compaction destroys soil structure and reduces rooting) and time of year. The soil infiltration rate and root zone depth should be used together to estimate the amount of water that needs to be available to the root system to avoid wilting. The rooting depth is multiplied by the PAW to estimate the total amount of water available to the turf.

Table 2. Available soil moisture and infiltration rates for common soil textures.^{1,2}

| Soil Texture | Soil Type | Typical plant-available moisture per foot of soil depth (inches) | Infiltration rate (inches h ⁻¹) |
|---------------|----------------------------|--|---|
| Light, sandy | Coarse sand | 0.7 | Fast (0.5 – 3) |
| | Fine sand | 0.9 | |
| Medium, loamy | Fine sandy loam | 1.5 | Moderate (0.25 – 0.5) |
| | Sil loam, loam | 1.9 | |
| Heavy, clay | Clay loam, Silty clay loam | 2.1 | Slow (0.1 – 0.25) |
| | Clay | 2.0 | |

3.7.3 Evapotranspiration

Evapotranspiration (ET) describes the water lost through soil evaporation and plant transpiration and is influenced by the climatic conditions on any given day, in addition to day length and solar intensity. For example, hot, windy days with low relative humidity have higher rates of ET than cooler, calm days with low relative humidity. Turfgrasses have crop coefficients of approximately 0.8 for cool season grasses and 0.6 for warm season grasses. To summarize, irrigating cool season grasses at 80% of weather station ET is used as the target for replenishing cool season turf water use so the plants do not experience water stress. An irrigation strategy

¹ Source: Adapted from Emmons, R.D. 2000. Turfgrass science and management, 3rd ed. Delmar Publishers, Albany, NY and published in [Drought Effects on Turf in the Landscape](#), UNL.

² Depth penetrated by 1” of water can vary depending upon a number of different soil factors. Therefore, this information is not included in the table.

during a prolonged drought, to conserve water, might be to practice deficit irrigation where 60 to 70 % of ET is applied. With that the turf might exhibit some browning from reduced irrigation.

Potential ET can be calculated and should be used to help determine irrigation needs. Sources of potential ET data include onsite weather stations and [National Weather Service](#) climate data.

3.7.4 Soil Moisture

Annual rainfall varies greatly in South Dakota. With the exception of the Black Hills, the southeastern section of the state receives the highest annual precipitation, and the northwest the lowest. The Black Hills receives nearly as much precipitation as the southeast. To accurately measure the local precipitation, the proper use of rain gauges, rain shut-off devices, soil moisture sensors, and other irrigation management devices should be incorporated into the site's irrigation schedule. Monitoring of soil moisture, in addition to calculating ET rates and visual observations of turf, assists in meeting turf water needs while conserving water resources.

3.7.5 Irrigation Scheduling

Proper irrigation can sustain plant energy reserves, increase root mass and depth, and reduce thatch accumulation. Irrigation should be applied as necessary to prevent wilt without oversaturating the soil/root zone and without compromising playing conditions. Soils do not need to be wetted below the root zone depth, as this could potentially lead to leaching, especially in shallow soils.

The goal of successful irrigation management is to limit excessive soil moisture while preventing wilt. Golf managers strive to precisely apply water so plant-available water is only slightly greater than predicted ET for any particular turfgrass. Applied irrigation is often greater than the measured weather station ET to compensate for irrigation distribution uniformities of less than 80%. For many highly maintained turf areas, like greens, small amounts of water are applied every night to replace what was lost the prior day. Soil moisture probes can help further improve irrigation precision. These technologies can guide irrigation head run times and identify locations that might benefit from additional hand watering.

During periods of sufficient natural precipitation, stress pre-conditioning with deficit irrigation can improve tolerance to future drought, heat, and cold stress. Deficit irrigation is the practice of limiting irrigation to slowly deplete soil PAW until the soil moisture approaches wilt points.

South Dakota experiences harsh, cold winters coupled with high wind velocities that may require the application of irrigation water during winter turf dormancy. During the summer, the turfgrass plant crowns are 80-85% water by weight. After fall cold-acclimation, crown moisture values typically range from 50-60% water by weight. This dramatically increases cold tolerance. When crown moisture values drop below 50% in winter, certain grass species, like creeping bentgrass, annual bluegrass, and perennial ryegrass, can suffer from desiccation stress. The lower the water weight (greater desiccation), the greater the risk of plant death when temperatures drop well below freezing.

Turf grown on sandy soils or having excessive thatch is more likely to have issues with winter desiccation. Therefore, lightly irrigating high value turf on dry, sunny days when the air temperatures are well above freezing is recommended to rehydrate plant crowns back to a survivable level and restore moisture at the surface. Other cultural practices such as sand topdressing, turfgrass covers, snow fences to collect snow cover, and desiccants may also help prevent desiccation/winter injury.

Computerized systems provide many advantages, including allowing a superintendent to remotely cancel the program if the course has received adequate rainfall. Clock-controlled irrigation systems preceded computer-controlled systems, and many are still in use today. Electric/mechanical time clocks cannot automatically adjust for changing ET rates. Therefore, frequent adjustment is necessary to compensate for the needs of individual turfgrass areas.

3.8 Irrigation Best Management Practices

Water Conservation and Efficient Use Planning

- ❖ Minimize acreage of irrigated turf.
- ❖ Convert out-of-play area turf to naturally adapted native plants, grasses, or ground covers to reduce water use and augment the site's aesthetic appeal.
- ❖ During times of intense heat stress, syringing, or the practice of applying a small amount of water to help cool the turfgrasses as it evaporates, may be beneficial under certain conditions.
- ❖ Create a drought management plan for the facility that identifies steps to be taken to reduce irrigation/water use and to protect critical and valuable areas.
- ❖ Adhere to local water use restrictions in time of drought. Use appropriate turfgrass species adapted to the location of the golf course and use drought-tolerant species whenever possible.

Irrigation Water Supply

- ❖ Identify appropriate water supply sources that meet seasonal and bulk water allocations for grow-in and routine maintenance needs.
- ❖ Use alternative water supplies/sources that are appropriate and sufficiently available to supplement water needs.
- ❖ Reclaimed, effluent, and other non-potable water supply mains must have a thorough cross-connection and backflow prevention device in place and operating correctly.
- ❖ Post signs in accordance with local utility and state requirements when reclaimed water is in use.
- ❖ Use salt-tolerant varieties of turf and plants to mitigate saline conditions, if necessary.
- ❖ Assess the irrigation water quality.
- ❖ Account for the nutrients in irrigation water when making fertilizer calculations.
- ❖ Monitor irrigation water regularly for dissolved salt content.
- ❖ Conduct a seasonal bulk water requirement analysis and a maximum bulk water requirement analysis.

- ❖ Design and/or maintain a system to meet a site's peak water requirements under normal conditions. Be flexible enough to adapt to various water demands and local restrictions.
- ❖ Install and maintain accurate metering systems.

Irrigation System Design

- ❖ Conduct a thorough site assessment prior to designing the irrigation system.
- ❖ Develop a written, site-specific Irrigation Management Plan.
- ❖ Seek assistance from irrigation professionals, such as from Certified Golf Course Irrigation System designers and [WaterSense-certified](#) irrigation consultants, and follow established BMPs related to system design.
- ❖ Sprinkler placement should avoid interfering with the playability of the hole.
- ❖ Irrigation pipes should be installed away from the green surface to avoid substantial increases in repairs and damages should pipe failures occur.
- ❖ Update multi-head control systems with single-head control systems to conserve water and to enhance efficiency.
- ❖ Manual quick-coupler valves should be installed for site specific irrigation, so these can be hand-watered during severe droughts.
- ❖ Install part-circle heads along lakes, ponds, wetlands margins, native areas, and tree trunks.
- ❖ Use part-circle or adjustable heads to avoid overspray of impervious areas such as roadways, sidewalks, and parking areas.
- ❖ Incorporate multiple nozzle configurations to add flexibility and enhance efficiency/distribution.
- ❖ When possible, use precise irrigation control technologies.

Irrigation System Installation

- ❖ The designer must approve any design changes before construction.
- ❖ Prior to construction, all underground cables, pipes, and other obstacles must be identified, and their locations flagged.
- ❖ Construction and materials must meet existing standards and criteria.
- ❖ Construction must be consistent with the design.
- ❖ Installers must provide an accurate and comprehensive As-Built map.

Irrigation System Maintenance and Performance

- ❖ Visual inspections should be conducted to identify necessary repairs or corrective actions, which should be completed before further evaluation of system performance.
- ❖ Pressure and flow should be evaluated to determine that the correct nozzles are being used and that the heads are performing according to the manufacturer's specifications.
- ❖ Pressure and flow rates should be checked (ideally annually) at each head to determine the average precipitation rates.
- ❖ Catch-can tests should be run to determine the uniformity of coverage and to accurately determine irrigation run times.

- ❖ Conduct an irrigation audit annually to facilitate a high-quality maintenance and scheduling program for the irrigation system.
- ❖ Inspect for water distribution interferences, such as trees and other obstructions.
- ❖ Inspect for broken and misaligned heads.
- ❖ Check that the rain sensor is present and functioning.
- ❖ Inspect the backflow device to determine that it is in place and in good repair.
- ❖ Examine turf quality and plant health for indications of irrigation malfunction or a need for scheduling adjustments.
- ❖ A professional irrigation audit that follows established guidelines is recommended at least every 8 years.
- ❖ Record any modifications to the As-Builts, including head and nozzle choices.
- ❖ Use photography to record and document any major underground installations/repairs.
- ❖ Review efficiency of above-ground electric motors annually.
- ❖ Licensed professionals should routinely inspect the well housing.

Irrigation System Winterization

- ❖ Flush and drain above-ground irrigation system components that could hold water as part of winter preparation.
- ❖ Remove water from all conveyances and supply and distribution devices that may freeze. Use compressed air or open the drain valves at the lowest point on the system.
- ❖ Change filters, screens, and housing; remove drain plugs and ensure any water is removed from the system. Secure systems and close and lock covers/compartments doors to protect the system from vandalism and from animals seeking refuge.
- ❖ Drain any above-ground pump casings that may have “trapped” water.
- ❖ Record metering data before closing the system.
- ❖ Secure or lock any remote irrigation components, including satellites.
- ❖ Perform pump and engine servicing/repair before winterizing.
- ❖ Recharge irrigation system in the spring with water and inspect for malfunctions.

Irrigation Management Decisions

- ❖ Evaluate root zone depth on the course and do not irrigate beyond this depth.
- ❖ Monitor potential ET and monitor plant available water to improve irrigation precision.
- ❖ Utilize soil moisture technologies and techniques consistently.
- ❖ Consider using soil moisture sensors to assist in scheduling or to create on-demand irrigation schedules.
- ❖ Consider using multiple soil moisture sensors to reflect soil moisture levels.
- ❖ Consider placing soil moisture sensors in a representative location within an irrigation zone.
- ❖ Use predictive models to estimate soil moisture and the best time to irrigate.
- ❖ Use a journal to record the “indicator zones” that should be more closely monitored.
- ❖ Calibrate older clock-control station timing devices periodically, and at least seasonally.

- ❖ Install control devices to allow for maximum system scheduling flexibility.
- ❖ Base plant water needs should be determined by ET rates, recent rainfall, recent temperature extremes, and soil moisture.
- ❖ Avoid use of a global setting; adjust watering times per head.
- ❖ Adjust irrigation run times based on current local meteorological data.
- ❖ Use a computed daily ET rate to adjust run times to meet the turf's moisture needs.
- ❖ Manually adjust automated ET data to reflect wet and dry areas on the course.
- ❖ Avoid irrigating beyond the point of runoff or leaching.
- ❖ Irrigation rates should not exceed the maximum ability of the soil to absorb and hold the water applied at any one time.
- ❖ Use infrequent, deep irrigation to supply sufficient water for plants and to encourage deep rooting in fairways and roughs.
- ❖ To maximize turf health during summer, irrigate to the depth of the turf root system in early morning.
- ❖ Visually monitor for localized dry conditions or hot spots to identify poor irrigation efficiency or a failed system device.
- ❖ The irrigation schedule should coincide with other cultural practices (for example, the application of nutrients, herbicides, or other chemicals).
- ❖ Respect the "water-in" requirement BMPs for fertilizer and plant protectant chemicals.

4 WATER MANAGEMENT

Whether natural or manmade, surface water storage in the form of lakes, ponds, and streams has long been associated with golf courses. Natural lakes and ponds are usually associated with existing water sources, such as wetland areas. Irrigation impoundments (lakes, ponds, and constructed wetlands) can be incorporated into the design of a course and used both to manage stormwater and to function as a source for irrigation.

South Dakota has abundant groundwater resources. For general information on aquifers, see the article [South Dakota Aquifers](#). Because groundwater is so plentiful, most of the state's citizens use it as drinking water. However, in some parts of the state, groundwater is contaminated, primarily with nitrates, other nutrients, and agricultural chemicals such as pesticides. Because of the public health concerns related to drinking water, preventing leaching into groundwater and protecting wellheads are important aspects of golf course management in South Dakota. Additionally, groundwater can be exchanged with streams. If there is negative aquifer pressure groundwater can become contaminated from polluted streams.

Overall, water management incorporates not only the information contained in this chapter, but many of the issues discussed throughout this document, including:

- Design considerations such as the use of vegetated buffers.
- Fertilization strategies near surface waters.
- Pesticide usage.
- Water quality monitoring.

In addition to managing stormwater to protect water quality and protecting groundwater resources, lakes and ponds on golf courses should be part of water management activities that deserve regular attention. Important parts of aquatic maintenance include managing components of aquatic habitats, such as algae and plant growth; reducing or preventing nutrient and sediment enrichment especially through the use of vegetated buffers; and ensuring dissolved oxygen levels needed to sustain aquatic life.

4.1 Regulatory Considerations

4.1.1 Water Right Permits

DENR is the state agency authorized to regulate surface and groundwater usage. All diversions of surface and groundwater for irrigation, hydropower, industrial use, municipal use, storage, and other uses require a [state water right permit](#). Reporting requirements are identified in permits. Superintendents should ensure that required reports are submitted on time.

4.1.2 Dams

DENR regulates the construction of dams in South Dakota for irrigation and requires a water right permit. DENR's [Building a Dam webpage](#) provides more information on the approval process.

4.1.3 Aquatic Pesticides

Aquatic pesticides that control nuisance aquatic plants like Eurasian milfoil, as well as algaecides that control algae, are available from commercial distributors. They require a [General Surface Water Discharge permit](#) from DENR. Some pesticides are listed as restricted use pesticide (RUP) and must be applied by a certified applicator. It is permissible for someone who is not a certified applicator to purchase or pick up a RUP with written permission from the certified applicator.

As with any pesticide application, the label must be followed. Labels on aquatic herbicides for algae control may specifically state that only a portion of the surface water area can be treated at a time to prevent massive algae and other plant die-offs and the low dissolved oxygen (DO) conditions that result from decaying organic matter.

4.1.4 Grass Carp

Though not prohibited in South Dakota, using grass carp (also known as white amur) for aquatic plant control is generally not a recommended practice. For more information on the disadvantages of stocking grass carp, see [Resolving Common Maintenance Problems /Aquatic Vegetation](#).

4.2 Stormwater Management

Best practices related to protecting the water quality of surface waters center on preventing nutrients, chemicals, and sediments from reaching waterbodies and wetlands. By managing stormwater effectively, maintaining buffers, and considering the special needs of wetlands, floodplains, lakes, and ponds, superintendents can effectively protect South Dakota's water resources.

The control of stormwater on a golf course is more than just preventing the flooding of the clubhouse, maintenance sites, and play areas. Proper management of stormwater controls the amount and rate of water leaving the course, controls erosion and sedimentation, stores irrigation water, removes waterborne pollutants, enhances wildlife habitat, and addresses aesthetic and playability concerns. Stormwater runoff (also called surface runoff) is the conveying force behind what is called nonpoint source pollution. Nonpoint source pollution is caused by water moving over and through the ground, picking up and carrying away natural and human-made pollutants, and finally depositing them into surface waters (lakes, rivers, wetlands, coastal waters) and groundwater. On golf courses, pollutants that might be found in surface runoff include, but are not limited to, pesticides, fertilizers, sediment, and petroleum.

Treating stormwater to avoid impacts to water quality is best accomplished by a treatment train approach in which water is conveyed from one treatment to another by conveyances that themselves contribute to the treatment. These treatments include source controls, structural controls, and non-structural controls. Source controls are the first car of the BMP treatment train. They help prevent the generation of stormwater runoff or the introduction of pollutants into stormwater runoff. The most effective method of stormwater treatment is to prevent or preclude the possibility of movement of sediment, nutrients, or pesticides in runoff.

The next car in the treatment system is often structural controls, which are design and engineering features of the course created to remove, filter, detain, or reroute potential contaminants carried in surface runoff. Examples of structural BMPs include ponds, constructed wetlands, and filters to address water quality, water recharge, and stream channel protection. Non-structural controls mimic natural hydrology and minimize the generation of excess stormwater and include vegetated systems. Vegetated systems such as stream buffers act as natural biofilters, reducing stormwater flow, removing sediments from surface water runoff, and preventing nutrient and pesticide discharge in runoff from reaching surface waters. The treatment train approach combines these controls, as in the following example: Stormwater can be directed across vegetated filter strips (such as turfgrass), through a swale into a wet detention pond, and then out through another swale to a constructed wetland system.



Figure 13. Retention ponds temporarily hold water from storm events.



Figure 14. Buffer strips, such as this one at Elkhorn Ridge Golf Club, are one of the most effective means of protecting surface water quality.

During any construction or redesign activity, proper erosion and sedimentation control must be followed (as discussed in the "Planning, Design, and Construction" chapter) to ensure that stormwater runoff does not impact water quality. Properly designed golf courses capture rain and runoff in water hazards and stormwater ponds, providing most or all of the supplemental water necessary under normal conditions, though backup sources may be needed during drought conditions.

4.2.1 Buffers

Buffers around the shore of surface waters, wetlands, or other sensitive areas filter runoff as it passes across the buffer. Buffers can be vegetated filter strips, such as those used as part of a stormwater treatment system. When used as a buffer along shorelines, stream banks, and wetland boundaries, filter strips are the last line of defense to keep sediment out of streams and to filter out fertilizers and pesticides that might otherwise reach waterways. Buffers also provide valuable food, cover, and travel corridors for some wildlife. Buffer widths as narrow as 15 feet have been shown to be effective in protecting water quality.

Care needs to be taken when applying pesticides or fertilizers near or in buffer strips to prevent movement into the nearby surface water. A number of turf species should be considered for buffer strips that can tolerate low inputs to avoid having to apply pesticides or fertilizers and require little to no mowing. [Research to evaluate low input grass species](#) in the north central area of the U.S., including South Dakota, shows that some species (tufted hairgrass, meadow fescue, blue grama, and prairie junegrass) could potentially provide turfgrass managers with a living turf stand when managed under low-input conditions and that some species (sheep fescue, hard fescue, and meadow fescue) might be able to survive under no-mow conditions. Evaluations were based on the following factors: lack of weed pressure, freedom from disease, plot density, and lack of lodging.

Riparian buffers along streams and rivers can be up to three different plant assemblages, progressing from sedges and rushes along the water's edge to upland species. Riparian buffers of sufficient width intercept sediment, nutrients, and pesticides in surface runoff and reduce nutrients and other contaminants in shallow subsurface water flow. Woody vegetation in buffers provides food and cover for wildlife, stabilizes stream banks, and slows out-of-bank flood flows. For more information, see the SDDA publication [Riparian Area Management](#).

4.3 Wetlands

The biological activity of plants, fish, animals, insects, and especially bacteria and fungi in a healthy, diverse wetland is the recycling factory of our ecosystem. Wetlands should be maintained as preserves and separated from managed turf areas by means of native vegetation, structural controls to protect water quality, and low/no maintenance activities to avoid nutrient or pesticide contamination.



Figure 15. Wetland and pond at Spearfish Canyon Golf Club; water quality is protected by a buffer strip around the perimeter..

4.4 Floodplains

Control structures located near floodplains, such as retention basins, store water and thereby reduce flooding and protect stream banks. These structures should be regularly inspected and maintained to ensure their proper function. Vegetated buffers along floodplains should be maintained to mitigate flooding, control stormwater, and protect water quality.

4.5 Lakes and Ponds

The management of lakes and ponds should include a clear statement of goals and priorities to guide the development of the BMPs necessary to meet those goals. Some of the particular issues superintendents should address to maintain the water quality of golf course lakes and ponds include:

- Pond design.
- DO (dissolved oxygen) levels.
- Aquatic plant management.
- Near-shore management zones.

4.5.1 Pond Location and Design

Designing a new pond requires considerations such as the size of the drainage area, water supply, soil types, and water depth. In addition to potentially serving as an irrigation water source, ponds support aquatic life and the construction of ponds should consider the needs of [aquatic ecosystems](#) (such as discouraging excessive growth of aquatic vegetation, DO needs for aquatic species, etc.). Careful design may significantly reduce future operating expenses for pond and aquatic plant management.

4.5.2 Dissolved Oxygen

Dissolved oxygen is the amount of oxygen present in water and is measured in milligrams per liter (mg/L). Adequate DO levels are required to sustain life in aquatic organisms and vary by species, the organism's life stage, and water temperature.

The amount of DO that water can hold depends on the physical conditions of the body of water (water temperature, rate of flow, oxygen mixing, etc.) and photosynthetic activity. Colder water has higher DO levels than warmer water. Dissolved oxygen levels also differ by time of day and by season as water temperatures fluctuate. Similarly, a difference in DO levels may occur at different depths in deeper surface waters if the water stratifies into thermal layers. Fast-flowing streams hold more oxygen than impounded water. Lastly, photosynthetic activity also influences DO levels. As aquatic plants and algae photosynthesize during the day, they release oxygen. At night, photosynthesis slows down considerably or even stops, and algae and plants pull oxygen from the water. In impoundments with excessive plant and algae growth, several cloudy days in a row can increase the potential for fish kills due to low DO during warm weather. Therefore, preventing excessive aquatic growth helps to maintain DO levels. The use of artificial aeration (diffusers) can also be used to maintain adequate DO, especially in small impoundments or ponds.

4.5.3 Aquatic Plants

Aquatic plants include algae and vascular plants. Phytoplankton, or algae, give water its green appearance and provide the base for the food chain in ponds. Tiny animals called zooplankton use phytoplankton as a food source. Large aquatic plants (aquatic macrophytes) can grow rooted to the bottom and supported by the water (submersed plants), rooted to the bottom or shoreline and extended above the water surface (emerged plants), rooted to the bottom with their leaves floating on the water surface (floating-leaved plants), or free-floating on the water surface (floating plants).

Aquatic plants are part of aquatic ecosystems. They provide a number of benefits, such as:

- Habitat for aquatic organisms (e.g. food and nesting sites).
- Oxygenation.
- Shoreline stabilization.

- Aesthetic appeal.

Aquatic plants growing on a littoral shelf may help protect receiving waters from the pollutants present in surface water runoff. Ideally, littoral zones should have a slope of about 1 foot vertical to 6-10 foot horizontal to provide the best substrate for aquatic plant growth. In open areas, floating-leaved and free-floating plants suppress phytoplankton because they absorb nutrients from the pond water and create shade.

Particularly in shallow or nutrient-enriched ponds, aquatic plant growth can become excessive. Non-native plants, in particular, can aggressively colonize aquatic environments. The excessive growth of any aquatic plant requires management. Following the principles laid out in the “Integrated Pest Management” chapter of this document, a number of controls should be considered to deal with excessive aquatic plant growth, including:

- Prevention, such as reducing nutrient (nitrogen and phosphorus) enrichment and avoiding the introduction of invasive species.
- Cultural practices, such as benthic barriers to prevent vascular plant growth.
- Mechanical removal.
- Chemical control.

Although grass carp are a biological control sometimes used to control aquatic plants, the use of this exotic species is not recommended, as noted in the "Regulatory Considerations" section earlier in this chapter.

For more on aquatic plant and algae management, see [*Resolving Common Maintenance Problems / Aquatic Vegetation*](#). Nebraska Game and Parks Commission.



Figure 16. Algae growth in an irrigation pond.



Figure 17. Aeration improves dissolved oxygen levels in ponds.

4.5.4 Shoreline Management

Special management zones should be established around the edges of lakes and ponds. The management specifications should include a setback distance when applying fertilizers, as well as reduced mowing. Grass clippings should be collected, as the phosphorus and nitrogen in clippings can impact the water quality.

4.5.5 Waterfowl

The deposits of fecal matter by resident and migrating waterfowl (such as Canada geese) can substantially impact water quality through nutrient enrichment. On golf courses, shallow ponds with significant populations of waterfowl are most likely to be affected. In addition, large numbers of Canada geese can erode shorelines and thin the grass cover on greens and fairways, contributing to the potential for erosion. Efforts to control waterfowl have met with mixed success. Loud sounds, dogs, and hunting have been tried in order to deter them. See the [South Dakota Games and Parks Goose web page](#) for more information on hunting geese in the state. However, many of these efforts do not lend themselves to golf courses, especially in more urban areas. For more information, see [Managing Canada Geese](#), University of Nebraska-Lincoln Extension.

4.6 Groundwater Management

Establishing protection zones around water supply wells and safe land-use practices that prevent leaching protect aquifers from accidental contamination.

4.6.1 Preventing Leaching

Leaching refers to the loss of water-soluble plant nutrients or chemicals from the soil as water moves through the soil profile and reaches the saturated zone. Some of the factors that can influence leaching potential include the depth to groundwater, soil type and structure, geology, rate of precipitation, and amount of irrigation. When applying fertilizers or pesticides, the rate, timing, and location of applications should be considered to minimize the potential for losses due to leaching. Sandy soils, for example, have a low potential to fix phosphorus and therefore are more likely to leach phosphorus. Nitrogen, in the form of nitrate (NO₃-N) presents leaching concerns for groundwater quality. Fertilizers with a solubility of more than 30 mg/L (or 30 ppm) can pose a risk for leaching.

4.6.2 Protecting Wellheads

The state's wellhead protection program allows municipalities, rural-water systems, or other governmental entities to establish land-use practices that will be allowed in the area of their water-supply wells or in the recharge area of their water supply. Protecting wellheads from physical impacts and contaminants, keeping them secure, and sampling wells according to the monitoring schedule required by the regulating authority are all best practices for ensuring drinking water is adequately protected. Before installing new wells, the DENR permit should be reviewed to determine the permitting and any setback requirements. New wells should be located up-gradient as far as possible from potential pollutant sources, such as petroleum storage tanks, septic tanks, chemical mixing areas, and fertilizer storage facilities. Most pesticide labels now prohibit mixing/loading pesticides within 50 feet (or other specified setback distance) from any well. Licensed water-well contractors are needed to drill new wells, which must meet regulatory and code requirements.

4.7 Water Management Best Management Practices

Stormwater Management

- ❖ Design stormwater treatment trains.
- ❖ Install berms and vegetated swales to capture pollutants and sediments from runoff before it enters irrigation storage ponds or other surface waters.
- ❖ Implement no- or low-maintenance vegetated buffer strips around surface waters.
- ❖ Utilize vegetated filter strips in conjunction with water filtration basins.
- ❖ Eliminate or minimize directly connected impervious areas.
- ❖ Use depressed landscape islands in parking lots to catch and filter water and allow for infiltration. When hard rains occur, an elevated stormwater drain inlet allows the island to

hold the treatment volume and settle out sediment, while allowing the overflow to drain away.

- ❖ When possible, maximize the use of pervious pavements, such as brick or concrete pavers separated by sand and plants.

Wetlands

- ❖ Maintain appropriate silt fencing on projects upstream to prevent erosion and sedimentation.
- ❖ Natural waters cannot be considered treatment systems and must be protected. (Natural waters do not include treatment wetlands.)
- ❖ Establish a low- to no-maintenance level within a 75-foot buffer along wetlands.
- ❖ Establish and maintain a 100-foot riparian buffer around wetlands, springs, and spring runs.

Floodplains

- ❖ Maintain stream buffers to restore natural water flows and flooding controls.
- ❖ Install buffers in play areas to stabilize and restore natural areas that also attract wildlife species.
- ❖ Install detention basins to store water and reduce flooding at peak flows.

Lake and Pond

- ❖ Maintain an unmowed, vegetated buffer strip to filter the nutrients and sediment in runoff, mowing only once or twice a year so that grasses and plants grow knee-high.
- ❖ If mowing near a pond or lake, collect clippings or direct them to upland areas so they do not increase nutrient loading to waterbodies.
- ❖ Maintain the required setback distance when applying fertilizers near waterways.
- ❖ Encourage clumps of native emergent vegetation at the shoreline.
- ❖ Maintain water flow through lakes if they are interconnected.
- ❖ Establish wetlands where water enters lakes to slow water flow and trap sediments.
- ❖ Maintain appropriate erosion and sedimentation controls on projects upstream to prevent sedimentation and nutrient enrichment to waterbodies.
- ❖ Dredge or remove sediment before it becomes a problem.

Dissolved Oxygen

- ❖ Establish DO thresholds to prevent fish kills, which occur at levels of 2-3 mg/L.
- ❖ Reduce stress on fish by keeping DO levels above 5 mg/L.
- ❖ Manipulate water levels to prevent low levels that result in warmer temperatures and lowered DO levels.
- ❖ Use artificial aeration (diffusers), if needed, to maintain adequate DO.

Aquatic Algae and Plants Management

- ❖ Develop a comprehensive management plan that includes strategies to prevent and control the growth of nuisance aquatic vegetation.
- ❖ Keep phosphorus rich material (e.g. natural or synthetic fertilizers, organic tissues like grass clippings, or unprotected topsoil) from entering surface water.

- ❖ Install desirable native plants to naturally buffer DO loss and fluctuation.
- ❖ To control excessive aquatic plant growth, use an IPM approach that incorporates prevention, cultural practices, and mechanical removal methods in addition to chemical control.
- ❖ To reduce the risk of DO depletion, use an algaecide containing hydrogen peroxide instead of one with copper or endothall.
- ❖ Dredge or remove sediment as needed to improve aquatic habitat.
- ❖ Reverse-grade around the waterbody perimeters to control surface water runoff and to reduce nutrient loads.
- ❖ Discourage large numbers of waterfowl from colonizing golf course waterbodies.
- ❖ Use a multi-faceted, IPM approach to control nuisance animals.

Preventing Leaching

- ❖ Identify areas on the course that may be prone to leaching (shallow depth to groundwater, sandy soils, etc.)
- ❖ Manage irrigation to avoid over-watering.
- ❖ Consider the potential for fertilizers or pesticides to leach before applying locations susceptible to leaching.

Wellhead Protection

- ❖ Use backflow-prevention devices at the wellhead, on hoses, and at the pesticide mix/load station to prevent contamination of water sources.
- ❖ Follow pesticide labels for setback distance requirements (typically a minimum of 50 feet).
- ❖ Properly decommission illegal, abandoned, or flowing wells.
- ❖ Surround new wells with bollards or a physical barrier to prevent impacts to the wellhead.
- ❖ Inspect wellheads and the well casing routinely for leaks or cracks; make repairs as needed.
- ❖ Maintain records of new well construction and modifications to existing wells.
- ❖ Obtain a copy of the well log for each well to determine the local geology and the well's depth; these factors will have a bearing on how vulnerable the well is to contamination.

5 WATER QUALITY MONITORING

Aligning golf course management practices with BMPs protects water quality on and downstream from the facility. A water quality monitoring program can confirm the effectiveness of a BMP-based program and provide important feedback on areas needing improvement.

Golf course superintendents seeking to develop and implement a monitoring program to document water quality conditions should first review available baseline water quality data. Baseline data can be assessed to determine the likely origin of contaminants, measure the extent of sedimentation and nutrient inputs, and estimate the potential impacts to surface water and groundwater. Following implementation of BMPs, routine monitoring can be used to measure water quality improvements and identify any areas where corrective actions should be taken.

Water quality monitoring on golf courses is voluntary. However, monitoring results demonstrate a commitment to water quality and implementing BMPs that protect water resources downstream. Furthermore, providing monitoring information to local, regional, and state regulatory authorities and watershed groups can help foster positive relationships with these stakeholders.

5.1 Regulatory Considerations

The EPA administers the protection of streams and lakes under the CWA. At the same time, the DENR creates state-specific regulations and water quality standards based on federal recommendations. Surface water quality is regulated under the CWA. DENR is the state's lead agency with regulatory authority for surface and groundwater quality. Surface water monitoring on golf courses is not a regulatory requirement, but it does demonstrate to regulators and the interested public the role of golf course superintendents in protecting the state's natural resources. The results of any monitoring programs should be compared with [South Dakota Surface Water Quality Standards](#).

The CWA requires states to prepare a list of impaired surface waters every other year. Impaired waters are those that do not meet the state water quality standards. From this list of impaired waters, states prepare Total Maximum Daily Loads (TMDLs) that include pollution control goals and strategies necessary to improve the quality of impaired waters and remove the identified impairments. TMDLs can include goals for nutrient loading (e.g. nitrogen or phosphorus). The DENR provides a list of finalized [TMDLs in the state](#).

In addition to developing TMDLs, DENR is required to provide Congress with surface water quality reports every two years that describe the status and trends of existing quality of all waters in the state. The report also provides information about the extent to which designated uses are supported. DENR combines this report with the impaired waters report into one [integrated report](#).

5.1 Existing Water Quality Information

Several sources of existing surface water and groundwater monitoring data may be available that can prompt a course-based water quality monitoring program. These potential data sources include:

- DENR's [Water Quality Monitoring Network](#).
- [South Dakota's Water Development Districts](#).
- EPA's [How's My Waterway](#) app.
- U.S. Geological Survey [water data for South Dakota](#).

5.2 Developing a Water Quality Monitoring Program

A water quality program begins with the development of a monitoring plan. The plan should identify specific conditions such as the presence of a watershed, stream flows, soil type, topography, drainage, and vegetation. In addition, the plan needs to document the hydrologic conditions and drainage, monitoring objectives, monitoring locations and frequency, and monitoring parameters. Baseline reference conditions can be established by collecting upstream water samples and comparing them with collection sites downstream of the areas influenced by golf course management practices.

Surface water collection sites can include streams, rivers, ponds, wetlands, etc., with the number and location of collection sites dependent upon the monitoring objectives. For example, a simple monitoring program can consist of the collection of DO data in surface waterbodies to ensure that these waterbodies can support aquatic life. Regardless of the extent of the monitoring program, the location of monitoring sites should remain consistent over time to establish trends in data. A more comprehensive monitoring program should include both field measurements at the time of sampling and analytical testing. Field measurements include pH, temperature, specific conductance, and DO. Lab testing should be conducted by a certified laboratory. Typical testing parameters include nutrients (such as nitrates and phosphorus), total dissolved solids (TDS), alkalinity, sediments, and selected pesticides used on the course. For more information on surface water monitoring programs, sampling procedures, and parameters specific to golf turf, the [Environmental Stewardship Guidelines](#) for Oregon courses can provide detailed guidance.

Developing a water quality monitoring program on golf courses is often limited to surface water monitoring. Sampling of stream macrobenthic invertebrates is a useful addition to a monitoring program, as macrobenthic species composition and diversity can be used as a relative assessment tool for stream health. For more information on macrobenthic invertebrate sampling, see New Mexico State University's [Stream Biomonitoring Using Macrobenthic Invertebrates](#). Such sampling can often be undertaken by university students in fulfillment of course work, by watershed association volunteer groups, or by other volunteer monitoring efforts.

In some instances, groundwater monitoring may be desired. Groundwater monitoring from wells located at the hydrologic entrance and exit from the course may be the best way to evaluate a

golf course's impact on water quality. If groundwater monitoring data from these locations are not available from existing sources, monitoring wells at the hydrologic entrance and exit from the course can be installed by private installers. Groundwater quality parameters can be limited to test only the ones directly influenced by course management, such as levels of pesticides and organic and inorganic nitrogen.

Water quality monitoring of irrigation sources (particularly water supply wells and storage lakes) provides valuable agronomic information that can influence nutrient programs. Immunoassay analysis may be a possible and cost-effective method for monitoring, depending on the analytical goals and the number of samples. To save money, several golf courses could pool resources and share immunoassay analyzer equipment and kits. See the "Irrigation" chapter of this document for more information on irrigation water quality issues.

5.3 Interpreting Water Quality Testing Results

Interpretation and use of water quality monitoring data depends to a large extent on the goal of the monitoring program. For example, the results may be analyzed to compare:

- Values over time.
- Values following implementation of BMPs, such as IPM measures.
- Monitoring points entering the site and leaving the site.

Results should also be interpreted and compared with the state's water quality standards, if water quality standards have been established for the parameter being evaluated. Data analysis can also be used to identify issues that may need corrective action, based on findings such as a spike in nutrient levels. For example, operator error in nutrient applications, an extreme weather event, or some combination of factors may be responsible. Water quality problems can often be addressed by simple changes to a course's existing nutrient management program.

5.4 Water Quality Monitoring Best Management Practices

Developing a Water Quality Monitoring Program

- ❖ Review existing sources of groundwater and surface water quality information.
- ❖ Develop a water quality monitoring program.
- ❖ Establish baseline quality levels for water.
- ❖ Identify appropriate sampling locations and sample at the same locations in the future.
- ❖ Visually monitor/assess any specific changes of surface waterbodies.
- ❖ Follow recommended sample collection and analytical procedures.
- ❖ Conduct seasonal water quality sampling. The recommendation is four times per year.
- ❖ Partner with other groups or volunteer water quality monitoring programs if possible, to share data and monitoring costs.

Interpreting Water Quality Testing Results

- ❖ Compare water quality monitoring results to benchmark quality standards.
- ❖ Use corrective measures when necessary.

6 NUTRIENT MANAGEMENT

Proper nutrient management plays a key role in the reduction of environmental risk and also increases course profitability. Among other benefits, applied nutrients increase the available pool of nutrients and allow turfgrass to recover from damage, increase its resistance to stress, and increase its playability. However, the increase in available nutrients also increases the potential risk of environmental impact. Nutrients may move beyond the turfgrass via leaching or runoff, which may directly impact water quality. Other organisms also respond to increases in nutrients and, in some cases, these organisms may deleteriously alter our ecosystem. The goal of a proper nutrient management plan should be to apply the minimum necessary nutrients to achieve an acceptable playing surface and apply these nutrients in a manner that maximizes their plant uptake.

6.1 Fertilizers Used in Golf Course Management

Understanding the components of fertilizers, the fertilizer label, and the function of each element within the plant are all essential in the development of an efficient nutrient management program.

6.1.1 Terminology

Grade or analysis is the percent by weight of nitrogen (N), phosphorous (P), and potassium (K) that is guaranteed to be in the fertilizer at minimum. Complete fertilizers contain N, P, and K.

6.1.2 Label

The label is intended to inform the user about the contents of the fertilizer. When applied according to the label, the use of fertilizer presents little to no environmental risk. Fertilizer labels generally provide the following information:

- Manufacturer's name and address.
- Brand name.
- Website.
- Nutrient guarantee (i.e. guaranteed minimum amounts of nutrients, given as a ratio).

Additional information that may be found on the label include characteristics such as size guide number (SGN), water-insoluble nitrogen (WIN), water-soluble nitrogen (WSN), and release characteristics.

6.1.3 Macronutrients

Macronutrients are required in the greatest quantities. N, P, and K are the ones most likely to be deficient in agronomic soils. Secondary macronutrients (calcium, magnesium, and sulfur) are also needed in relatively high quantities but are rarely deficient in turf soils. Understanding the role of each macronutrient within the plant provides a greater understanding of why these nutrients play such a key role in proper turfgrass management.

The Role of N

Nitrogen is required by the plant in greater quantities than any other element except carbon, hydrogen, and oxygen. Nitrogen plays a role in numerous plant functions and is an essential component of amino acids, proteins, and nucleic acids.

The goal of all applied nutrients is to maximize plant uptake while minimizing nutrient losses. Understanding each process below, in addition to the fate and transport mechanisms of runoff and leaching, leads to sound decision making and ultimately leads to a reduction in environmental risk. These nitrogen processes are:

- Mineralization, the microbial mediated conversion of organic N into plant-available NH_4 .
- Nitrification, the microbial-mediated conversion of NH_4 to NO_3 .
- Denitrification, the microbial mediated conversion of NO_3 to N gas; this primarily occurs in low-oxygen environments and is enhanced by high soil pH.
- Volatilization, the conversion of NH_4 to NH_3 gas.

Understanding how certain N sources should be blended and applied is an essential component in an efficient nutrient management plan. In many cases, N sources are applied without regard to their release characteristics. This is an improper practice and increases the risk of negative environmental impact. Each N source (particularly a slow-release form) is unique and therefore should be managed accordingly. For example, applying a polymer-coated urea in the same manner as a sulfur-coated urea greatly reduces the value of the polymer-coated urea. Similarly, applying 2 pounds of N from ammonium sulfate may cause burning, while applying 2 pounds of N from certain polymer-coated ureas may not provide the desired turfgrass response. Rate, application date, location, and turfgrass species all should be included in nutrient application decision-making.

Soluble Nitrogen Sources

Soluble N sources include:

- Urea (46-0-0).
- Ammonium nitrate (34-0-0).
- Ammonium sulfate (21-0-0).

- Diammonium phosphate (18-46-0).
- Monoammonium phosphate (11-52-0).
- Calcium nitrate (15.5-0-0).
- Potassium nitrate (13-0-44).

Slow-release Nitrogen Sources

A slow-release N source is any N-containing fertilizer where the release of N into the soil is delayed either by requiring microbial degradation of the N source, by coating the N substrate to delay the dissolution of N, or by reducing the water solubility of the N source. Slow-release nitrogen sources include:

- Coated urea (sulfur, polymer, or both).
- Urea-formaldehyde/urea-formaldehyde reaction products (methylene urea).
- Isobutylidene diurea (IBDU).
- Natural organic compost (plant material or animal manure and biosolids).

Urease and Nitrification Inhibitors

Urease inhibitors reduce the activity of the urease enzyme resulting in a reduction of volatilization and an increase in plant-available N. Nitrification inhibitors reduce the activity of *Nitrosomonas* bacteria, which converts NH_4 to NO_2 . This reduced activity results in a reduction of N lost via denitrification and an increase in plant-available N. These compounds are best applied to turfgrass sites that are not irrigated.

The Role of Phosphorous

Phosphorus forms high-energy compounds that are used to transfer energy within the plant. Phosphorus may remain in an inorganic form or may become incorporated into organic compounds.

Phosphorus can be a growth-limiting factor for many unintended organisms and is a major contributor to eutrophication of waterbodies. Therefore, the goal of P fertilization is to supply enough to sustain healthy turfgrass growth yet minimize environmental risk. Phosphorous application rates should be based on soil test results. Fortunately, phosphorus binds tightly to native soils and organic matter. The most important way to limit phosphorus movement off-site is to maintain dense and healthy vegetation.

P Fertilizer Sources

Phosphorus sources include:

- Diammonium phosphate.
- Concentrated superphosphate.
- Monoammonium phosphate.
- Potassium phosphate.
- Natural organics.

The Role of Potassium

Potassium is of no environmental concern, but can be an economic concern, particularly when potassium is overutilized, which can be quite common. Generally, potassium concentrations in turfgrass tissue are about one-third to half that of N. Potassium is not a component of any organic compound and moves readily within the plant. Potassium is a key component of osmoregulation and has been documented to increase stress resistance.

Potassium Fertilizer Sources

Potassium fertilizer sources include:

- Potassium chloride.
- Potassium nitrate.
- Potassium phosphate.
- Potassium sulfate.

6.1.4 Secondary Macronutrients

Fertilization with secondary macronutrients is rarely justified in a turfgrass systems. Calcium (Ca), magnesium (Mg), and sulfur (S) are essential to plant function, but fibrous turfgrass root systems and soils typically satisfy demand for these nutrients. Soil test interpretations for these nutrients are unsubstantiated in turf because deficiencies for these nutrients are exceedingly rare. Such deficiencies have only been created in hydroponic culture (no soil). Therefore, suspected secondary nutrient deficiencies should be confirmed with tissue tests. Also, confirm suspected deficiencies with small-test applications of fertilizer. If there is not a turfgrass response, then a widespread nutrient application is not warranted. Secondary nutrient sources can be found in Table 3.

Table 3. Secondary macronutrients

| Nutrient | Role | Sources |
|-----------|--|---|
| Calcium | Primarily a component of cells walls and structure. | Gypsum Limestone Calcium chloride |
| Magnesium | Central ion in the chlorophyll molecule and chlorophyll synthesis. | S-Po-Mg Dolomitic limestone Magnesium sulfate |
| Sulfur | Metabolized into the amino acid cysteine, which is used in various proteins and enzymes. | Ammonium sulfate Elemental sulfur Gypsum Potassium sulfate |

6.1.5 Micronutrients

Micronutrients are just as essential for proper turfgrass health as macronutrients but are required in very small quantities compared with macronutrients. Micronutrients include iron (Fe), manganese (Mn), boron, copper, zinc, molybdenum, and chlorine. They play a variety of roles in turf biology, including roles in photosynthesis, nitrogen fixation, protein synthesis, etc. Micronutrient deficiencies can be confirmed by tissue testing or small fertilizer applications to turf to verify fertilizer response. Soil testing for micronutrients is not recommended, and soil interpretations for these nutrients can be ignored.

Iron and manganese deficiency symptoms can be common in bluegrasses and bentgrass during summer. Deficiency symptoms include yellow colored (chlorotic) turf that does not respond to nitrogen fertilization. In many instances, nitrogen fertilization will intensify the chlorosis. The chlorosis is most severe when soils are warm, wet, and have high pH (>7.3). It is believed that root function is lost under these conditions. As a result, foliar iron and manganese applications will effectively correct the deficiency. Deficiency symptoms subside as the soil cools into the fall.

6.2 Soil pH

Soil pH influences nutrient availability and microbial communities. Unlike some pH-sensitive plant species, turfgrasses are largely pH-independent. The dense and fibrous root system of turfgrasses allow turf plants to thrive across a broad pH range. Soil pH ranging from 5.5 to 8.0 provides acceptable nutrient availability for most turf stands.

Application of lime following soil test recommendations may be recommended if soil pH is less than 5.5 to optimize nutrient availability and reduce the risk of aluminum toxicity. Lowering pH is rarely successful in South Dakota because many of the soils are highly buffered with calcium carbonate (limestone) parent material. This is typically true if the soil pH is >7.3. Note that sensitive species may benefit from iron and manganese fertilizer during the summer in these high pH soils.

6.3 Soil Testing

Soil tests estimate nutrient availability and predict a plant's response to an applied nutrient. Accurate and consistent sampling and analysis provide useful soil test information over time. Soil test sampling and analytical testing recommendations include the following:

- Conduct soil sampling at a 4-6" depth from representative areas of similar management.
- Exclusively use one trusted soil testing laboratory.

Soil testing is best for providing guidance for soil pH, organic matter content, plant available phosphorus and potassium, and salinity and soil sodicity. Soil tests results and interpretations are provided in Table 4. Soil test result recommendations for other nutrients are not supported by turf science. The best way to utilize soil testing is to monitor changes in the nutrients over time. Declining soil test phosphorus and potassium, for example, indicates that fertilization with P₂O₅ and K₂O should be increased relative to nitrogen fertilization rate, unless the values for those nutrients are deemed to be high. Nitrogen fertilizer drives the uptake of all other nutrients in turf. Therefore, P₂O₅ and K₂O should be looked at relative to N fertilization rate and not just the total amount applied.

For more information, see the following:

- [*Simplifying Soil Test Interpretations for Turf Professionals*](#), UNL Extensions.
- [Soil testing labs](#) information for South Dakota.

6.4 Nutrient Management Planning

Fertilizer programs are ultimately designed to supply nutrients to the turf as they diminish over time. For example, nutrients can leave a turf system through soil and water runoff, nutrient leaching, volatilization, and mowing. Nutrients can also be temporally removed from a system through processes of nutrient fixation and immobilization. Nitrogen is particularly difficult to manage because it can be quickly immobilized and then mineralized depending on a soil's physical, chemical, and biological properties. Therefore, the goal of a successful nutrient management program should be to sustain even levels of plant-available nutrients for a uniform growth rate and sustain adequate recuperative potential to meet expectations of quality and turf performance, while minimizing excessive growth and the risk of nutrient loss to the environment.

Record keeping and review can help to finetune the amount, timing, placement and rate of application of nutrients as prescribed by soil testing. Important records to keep for nutrient management planning include the following:

- Site maps
- Soil test results for greens, tees, fairways, roughs, and clubhouse grounds
- Fertilizer application records for each area

Table 4. Soil test results and recommendations

| Soil Test Result | Desired Value | Soil Test Result | Annual Fertilizer Recommendation |
|--|--|--|--|
| Soil pH | 5.5-8.0 ^c | Less than 5.5 | Consider lime application |
| | | 6.0-8.0 | No remediation required |
| | | Greater than 8.0 | Consider use of acidifying fertilizer; potential micronutrient limitation |
| Soil organic matter (SOM) | No recommended range | Much greater than previous year | Reduce inputs (nitrogen and water) Increase cultivation and topdressing |
| | | Slightly greater or same as previous year | Some increase is normal in new turf stands / continue good management |
| | | Much less than previous years | Likely the result of aggressive cultivation and/or reduced inputs |
| Phosphorus (P) | 25-50 ppm ^a | Less than 25 | >0.25 lb P ₂ O ₅ per lb N applied ^b |
| | | 25-50 | 0.25 lb P ₂ O ₅ per lb N applied ^b |
| | | Greater than 50 | No P fertilizer required |
| Potassium (K) | 40-80 ppm ^a | Less than 40 ppm | >1 lb K ₂ O per lb N applied ^b |
| | | 40-80 ppm | 0.75 to 1 lb K ₂ O per lb N applied ^b |
| | | Greater than 80 ppm | No K ₂ O required |
| Salinity | < F3 dS/m ^c | Less than 1.5 dS/m | Low salinity risk |
| | | 1.5 to 3.0 dS/m | Bluegrasses sensitive, leach soil |
| | | Greater than 3.0 dS/m | Most turfgrasses sensitive, leach soil |
| Sodicity (native soils only) | < 5% ESP | Less than 5% ESP | Low sodium risk in fine-texture soil |
| | | 5-15% ESP | Consider gypsum treatment to improve permeability of native soils |
| | | Greater than 15% | Sodic soil, treat native soils with gypsum |
| All other nutrients (Ca, Mg, S, N, Fe, etc.) | No science-based soil test interpretations | Confirm deficiency with tissue testing or small applications to turf to verify fertilizer response | All other nutrients (Ca, Mg, S, N, Fe, etc.) |

^a Mehlich-3 soil test method.

^b Demand for P and K fertilizer is affected by N fertilizer, soil type/environment, and clipping management. For example, turf on a native soil, clippings removed, and fertilized annually with 4 lbs of N/1,000 ft² would need about 1 lb of phosphorus (P₂O₅) and 3 lbs of potassium (K₂O)/1000 ft² to sustain soil test levels. Returning clippings reduces those P and K requirements by 50%. These ratios are good starting values and may need to be adjusted to sustain soil test P and K levels at any particular location.

^c Saturated soil paste extract method.

Nutrient management plans incorporating this information is an additional step that can help to inform golf course management activities.

6.4.1 Nitrogen Fertilization

Nitrogen is the most important nutrient managed by golf course superintendents as it drives turf growth rate. During the growing season, slow growth is an indication of low soil N status. Alternatively, excessive amounts of growth typically indicate there is large pool of available N. That fertilizer may have come from fertilizer application or organic N mineralization within the soil.

Research on a Kentucky bluegrass lawn shows that only 1 pound of the 3 pounds of N taken up during a year came from synthetic fertilizer. The remainder came from the soil or clippings.

Soil N mineralization is typically greatest right after winter (during spring greenup) and when the soil is warm and wet during mid-summer. During these periods, turfgrass growth rate is accelerated and N fertilization should be avoided. Soils with high levels of organic matter generally have more mineralization than newer turf stands with less organic matter. As a result, N fertilizer requirements will be higher in new turf stands than well-established turf stands. Those requirements will decline as the turf stand ages. The exact time depends on many management, use, and soil factors.

Optimum N fertilization times occur when soil nitrogen mineralization is low. This typically occurs in late May and into June and then from late August and September, across South Dakota. While mineralization is low in late fall and during winter, plant uptake is also diminished as it starts to gain cold-hardiness into and through winter. Nitrogen fertilizer should not be applied after mid-October because uptake is low and loss to the environment is greatest.

Monitor turfgrass performance, including color, growth rate, and traffic recovery to help schedule N application timings. Best cool-season turf application times are in late spring and early fall, following traditional spring and summer growth flushes. Cool-season turf stands can benefit from summer fertilizer application if mineralization is low and turf is underperforming. Warm season turf should be fertilized after greenup and again in mid to late summer. Late summer applications can help sustain green color of warm-season grasses into the fall.

6.4.2 Nitrogen Application Rates

Turfgrass is extremely responsive to N fertilizer. Highly maintained turf areas (greens and tees) are traditionally fertilized by “spoon-feeding” small amounts of N frequently. This approach has several advantages. First, the small quantity of N (typically sprayed on the turf) presents a smaller risk for a large leaching or other loss event. Light and frequent applications also increase control because managers can vary fertilizer rates and intervals based on the performance of the turf. Highly soluble sources are recommended for these applications because they are fast-acting and short-lived. Application rates when spoon-feeding with soluble N sources typically range from 0.0 to 0.5 lb of N per 1,000 square feet. Applications at rates above 0.75 lb of N per 1,000

square feet can increase the risk of fertilizer burn and N loss. In most cases, fertilizer should be watered-in with less than 0.2” of irrigation to minimize the risk of burn or nutrient loss.

Turf areas that are less intensively managed are traditionally treated with granular products, which can range from 100% water soluble products to 100% controlled-release products. Most application rates range from 0.5 to 2.0 lbs of N per 1,000 square feet (Table 5). Lower rates are more common for quick-release fertilizers and fertilizers with a small size guide number (SGN), the median prill diameter in millimeters x 100. Products with SGN < 100 are best for putting greens, 120-180 are best for fairways, and >200 are best for roughs. Higher N rates are most common with large SGN products and products that have very slow-release characteristics or large SGN materials. Some slow-release fertilizer products can release fertilizer over a period of 75 days to 120 days after application. An obvious advantage to these very slow release products is fewer applications. A disadvantage is the manager forfeits control; little can be done if the turf is growing too rapidly following an application.

The ideal time to fertilize cool-season species is late spring and early fall and warm-season species during the summer. Products with more water soluble (quick-release) N should be used as the season starts to end. All fertilizer N should have become available by the end of the growing season.

Table 5. Annual nitrogen application rates

| Turf managed area and species | Pounds of actual N / 1,000 sq. ft. annually* |
|-------------------------------|--|
| Greens | |
| Bentgrass | 2-4 |
| Annual bluegrass | 2-5 |
| Tees | |
| Creeping bentgrass | 1-5 |
| Annual bluegrass | 2-5 |
| Kentucky bluegrass | 2-5 |
| Perennial ryegrass | 2-6 |
| Buffalograss | 1-3 |
| Fairways | |
| Creeping bentgrass | 0-3 |
| Kentucky bluegrass | 1-3 |
| Perennial ryegrass | 2-5 |
| Buffalograss | 0-3 |
| Roughs | |
| Fine fescues | 0-2 |
| Kentucky bluegrass | 0-3 |
| Perennial ryegrass | 2-4 |
| Turf-type tall fescue | 0-2 |

| | |
|--------------|-----|
| Buffalograss | 0-3 |
|--------------|-----|

** Annual N fertilizer rates will be affected by soil organic matter, climate, length of growing season, cultural practices, traffic, and aesthetic expectations. Mature turf sites with little traffic may require little to zero N. However, new sites that are highly trafficked may require more than the values in this table.*

6.4.3 Phosphorus and Potassium Recommendations

The goal of P and K fertilization is to supply enough to sustain healthy turfgrass growth yet minimize unnecessary cost and environmental risk. Soil test calibration and interpretations suggest that soil test P levels should remain above 20 ppm Mehlich-3 P or between 15-20 using the Olsen method. Potassium levels should be greater than 40 ppm Mehlich-3 K. Apply P and K following recommendations in Table 4.

6.5 Fertilizer Applications

Because N and P are two of the major sources contributing to both surface and groundwater pollution, fertilizers must be applied considering the application site and prevailing conditions. For example, sandy soils often have a lower potential to fix P and are more likely to have a leaching problem than other soils. Fertilizer applications should be avoided whenever possible on steep slopes and should not be allowed to be deposited on impervious surfaces, such as paved cart paths and parking areas. To prevent both runoff and leaching, avoid applying fertilizer to soils that are at, or near, field capacity or following rain events that leave the soils wet. In addition, maintaining a vegetated buffer or filter strip around surface waterbodies significantly filters out any nutrients in surface runoff.

The selection and calibration of application equipment is an important aspect of nutrient management. Not all fertilizers can be spread with every spreader. For example, if sulfur-coated urea is spread through a drop spreader, the sulfur coating could be damaged, essentially leading to an application of soluble urea. Therefore, choosing the appropriate spreader for a given material (walk-behind rotary drop spreader, bulk rotary, or spray) is important.

Accurately calibrated sprayers or spreaders are essential for proper application of fertilizers. Incorrectly calibrated equipment can easily apply too little or too much fertilizer, resulting in damaged turf, excess cost, and greater potential of nutrient movement off-site. An excellent resource for spreader care and calibration can be found at [Penn State's Department of Plant Science](#). Spreaders should also be thoroughly cleaned after use due to the high salt content that corrodes metal parts and in keeping with the BMPs for equipment washing.

6.6 Nutrient Management Best Management Practices

- ❖ Keep accurate records of fertilizer applications.
- ❖ Because turf is extremely responsive to soil N status, evaluate changes in clipping yield during the growing season to estimate N availability.

- ❖ Reduce N inputs on more mature turfgrass stands.
- ❖ Use N fertilizer to produce an even growth rate. This increases golf course playability and minimizes the risk to the environment, while excessive fertilization reduces playability and increases the risk of N leaching.
- ❖ Evaluate the previous fertilizer application for response prior to a follow-up application.
- ❖ Use soluble N sources (0.05-0.50 lb N per 1,000 square feet) to fine-tune clipping yield on highly managed turf surfaces.
- ❖ Fertilizer products with a blend of quick and slow release fertilizer are frequently applied to non-intensively managed areas. Optimum timing for cool-season species are late spring/early summer and again in late summer/early fall.
- ❖ Summer fertilizer applications can benefit young turf stands or stands growing on poor soils.
- ❖ Apply fertilizer when turf is actively growing to minimize loss.
- ❖ Light irrigation after P application has been shown to reduce P runoff.
- ❖ Maintain dense turf stand through proper nitrogen fertilization to reduce the potential for soil runoff.
- ❖ Monitor P and K by testing soil regularly.

Fertilizer Applications

- ❖ Prevent fertilizers from being deposited onto impervious surfaces.
- ❖ Avoid applying fertilizer to soil at or near field capacity or following rain events that leave the soil wet.
- ❖ Do not apply fertilizer when heavy rains are likely.
- ❖ Maintain buffer areas around waterbodies. The buffer areas should not be not fertilized.
- ❖ Choose the appropriate type of spreader for a given fertilizer.
- ❖ Calibrate application equipment regularly.
- ❖ Maintain records of applied fertilizer rates in lbs/1000 sq. ft. per application
- ❖ Summarize annual nutrient amounts per growing season and adjust as needed for planning purposes the following year.

7 CULTURAL PRACTICES

Cultural practices make up cultural programs that vary by the distinct turfgrassed area on the golf course. Proper cultural management can help produce a dense, healthy turf playing surface. These practices are used on all areas of a golf course, including putting greens, fairways, tee boxes, and roughs and include a variety of methods, such as mowing, cultivation, cultivar selection, and rolling. These practices typically manage the top 3-4" of soil and improve plant nutrient/water uptake and the overall health of the plant.

7.1 Mowing

Mowing is the most commonly used cultural practice on golf courses. Mowing practices impact turf density, texture, color, root development, and wear tolerance. Failure to mow properly will result in weakened turf with poor density and quality. Mowing height decisions are typically based on the turf species and location on the course. Other factors affect mowing as well, such as frequency, shade, equipment, time of year, root growth, and abiotic and biotic stress. Mowing should increase tillering and shoot density while not decreasing root and rhizome growth as much as possible.

7.1.1 Height of Cut

Height of cut (HOC) is important for a healthy playing surface. Proper mowing HOC is a function of the species/cultivar being managed and the intended use of the site. While taller grown turf is more likely to withstand pests and stresses, a well-groomed dense turf stand is preferred by many golfers for playability and aesthetic appeal.

Various heights of cut are used on different locations on a golf course. Table 6 shows recommended HOC for turf species that will help to maximize turf density, assuming water, nutrient, and cultivation needs are being met. Follow the rule of thumb that no more than one-third of the plant should be removed at one mowing to avoid scalping, which reduces turf density and can result in a dramatic reduction in root growth.

Mowing heights can vary seasonally, due to such factors as sun availability, weather, and drought stress. During the spring and fall, when turf does not experience heat stress, certain mowing heights can affect the plants disease susceptibility. Throughout the heat of the summer, lower cuts can increase stress. Therefore, a higher HOC is recommended to help insulate the crown from heat stress, reduce weed competition, and reduce water needs. In shaded areas, photosynthetically active radiation is restricted, and the turfgrass response is to grow upright in an effort to capture more light to meet photosynthetic needs. To counterbalance this effect, mowing HOC should be increased for turf grown in shaded environments. Alternatively shade removal or control favors healthy turfgrass growth.

Table 6. Mowing HOC recommendations by species and location

| Turf Species | Greens | Tees, Collars, Approaches | Fairways | Roughs |
|--------------------|-------------|---------------------------|------------|---------|
| | (in inches) | | | |
| Creeping bentgrass | 0.10-0.16 | 0.2 – 0.6 | 0.25 – 0.5 | - |
| Kentucky bluegrass | - | 0.25 – 0.8 | 0.25 – 0.8 | 2 – 4 |
| Annual bluegrass | 0.10-0.16 | 0.2 – 0.6 | 0.25 – 0.5 | - |
| Perennial ryegrass | - | 0.4 – 0.6 | 0.4 – 0.6 | 2 – 4 |
| Tall fescue | - | - | - | 2 – 4 |
| Fineleaf Fescue | - | - | 0.6 – 0.8 | 2 – 4 |
| Buffalograss | - | - | 0.6 – 0.75 | 1.5 - 3 |

7.1.2 Frequency

Mowing frequency should be based on vertical leaf growth. Maintaining an optimal root-to-shoot ratio is critical. Turfgrass plants that are mowed too low will require a substantial amount of time to provide the food needed to produce shoot tissue for future photosynthesis. If turf is mowed too low in one event, an imbalance occurs between the remaining vegetative tissue and the root system, resulting in more roots being present than the plant needs physiologically. As a result, the plants will slough off the unneeded roots. Root growth is least affected when no more than 30% to 40% of leaf area is removed in a single mowing. Therefore, when lowering mowing heights the adjustments should be gradual to allow the turf to adjust to a new system. Mowing frequency should be decreased to allow the turf to recover when is under extremes, such as heat, drought, heavy traffic, increased mowing frequency, or other cultural practices (e.g. aeration).

7.1.3 Patterns

Mowing patterns influence both the aesthetic and functional characteristics of a turf surface. Changing the direction of cut is used to prevent excessive lateral growth and avoid lay over, providing a cleaner playing surface and an easier maintained HOC. While patterns should be varied regularly throughout the course, the direction of cut should be changed on putting greens every time it is mowed, including changing the direction of clean up and skipping some clean-up mows. Varying mowing patterns also provides aesthetic value.

7.1.4 Clipping Management

Turfgrass clippings are a source of nutrients, containing 2% to 4% nitrogen on a dry-weight basis, as well as significant amounts of phosphorus and potassium. These nutrients can be sources of pollution, and therefore should be handled properly to avoid contaminating water resources. Clippings are best returned to the site during the mowing process unless the presence of grass clippings will have a detrimental impact on play, such as on putting greens, or when the

amount of clippings is so large that it could smother the underlying grass. Collected clippings should be disposed of properly to prevent undesirable odors near play areas. Consider composting clippings or dispersing them evenly in natural areas where they can decompose naturally without accumulating in piles, though care should be taken to ensure that clippings are free from pesticides.

7.1.5 Mowing Equipment

Different mowing equipment is typically used on different locations of a golf course. For example, reel mowers are ideally suited for maintaining turfgrass stands that require a HOC below 1.5” and provide the best quality cut when compared with other types of mowers. Rotary mowers, when sharp and properly adjusted, deliver acceptable cutting quality for turf cut above 1” in height. Flail mowers are most often used to maintain infrequently mowed areas.

Maintaining blades by sharpening and adjusting them regularly provides the best quality cut. Dull blades shred leaf tissue, increase water loss, and increase the potential for disease development.

7.2 Aeration

Turfgrasses are the only plant materials in the landscape upon which we walk, drive golf carts, drive mowing equipment, and take divots with clubs and make indentations with golf balls on very closely mowed golf green. Traffic results in soil compaction and wear on the turfgrass leaves. Golf is a game that relies on managing smooth surfaces on greens tees and fairways. Cultivation practices -- aeration practices and surface cultivation practices -- disturb the soil or thatch through the use of various implements to achieve important agronomic goals that include relief of soil compaction, thatch/organic matter reduction, and improved water and air exchange. Aeration practices consist of core aeration, deep drilling, solid tining, and high-pressure water injection. Light and frequent sand topdressing applications are also beneficial for smoothing the surface, diluting organic matter, and improving playability.

Aeration frequency depends upon traffic intensity, thatch/organic matter buildup, black layer and level of soil compaction. Even though aeration is very beneficial, it disturbs the playing surface and takes some time to heal. Table 7 shows advantages and disadvantages of multiple aeration practices.

Table 7. Aeration practices

| Method | Compaction relief | Surface disruption | Water/air movement | Disruption of play |
|-------------------------------|--------------------------|---------------------------|---------------------------|---------------------------|
| Hollow-tine aeration | High | Medium | High | Medium to High |
| Deep drilling | Medium | Medium | High | High |
| Solid-tine aeration | Low to High | Low | High | None to Medium |
| High-pressure water injection | None | Low | High | None to Low |

7.2.1 Hollow-tine Aeration

Hollow-tine (or core) aeration is effective at relieving soil compaction, improving internal soil drainage and increasing oxygen in the soil. Aeration involves physically removing cores, varying in depth, diameter, and distance apart. Table 8 shows different core sizes used for aeration.

Table 8. Core size options for aeration.

| Tine Size (in.) | Spacing (in.) | Holes/ft ² | Surface Area of One Tine (in.) | Percent Surface Area Affected (Outside tine) |
|-----------------|---------------|-----------------------|--------------------------------|--|
| 1/4 | 1.252 | 100 | 0.049 | 3.4% |
| 1/4 | 2.52 | 25 | 0.049 | 0.9% |
| 1/2 | 1.252 | 100 | 0.196 | 13.6% |
| 1/2 | 2.52 | 25 | 0.196 | 3.4% |
| 5/8 | 2.52 | 25 | 0.307 | 5.3% |



Figure 18. Core aeration holes.



Figure 19. Fairway aeration.

7.2.2 Deep Drilling

Deep-drill aeration creates deep holes in the soil surface profile through use of drill bits. Soil is brought to the surface and distributed into the canopy. Holes can be backfilled with new root-zone materials if a drill-and-fill machine is used. These machines allow replacement of heavier soils with sand or other materials in an effort to improve water infiltration into the soil profile.

7.2.3 Solid-Tine Aeration

Solid tines cause less disturbance to the turf surface and can be used to temporarily reduce compaction and soften surface hardness during months when the growth rate of grasses is reduced. Benefits of solid-tine aeration are temporary because no soil is removed from the profile. Solid tining without sand topdressing will impact water infiltration but has little impact on organic matter management.

7.2.4 High-Pressure Water Injection

High-pressure water injection promotes water penetration and air exchange. Streams of water are injected at high velocities 4-8" into the soil at 1/8" to 1/4" diameter. High-pressure water injection doesn't disrupt play.

7.3 Surface Cultivation

The goals of surface cultivation are to manage excess organic matter accumulation above the soil, reduce the formation of leaf grain, improve infiltration, and improve surface consistency (Table 9). These methods are generally less disruptive than traditional aeration practices and can quickly impact a large percentage of the turfgrass canopy. They usually have low to no impact on soil compaction.

Table 9. Surface cultivation practices

| Method | Compaction relief | Surface disruption | Water/air movement | Disruption of play |
|-----------------|-------------------|--------------------|--------------------|--------------------|
| Vertical mowing | None to Low | Medium to High | Medium | Low to High |
| Spiking/slicing | None | Low | Low | None |

7.3.1 Vertical Mowing

Vertical mowing can be incorporated into a cultural management program to achieve a number of goals. The grain of a putting green can be reduced by setting a verticutter to a depth that just nicks the surface of the turf. Frequent, light vertical mowing minimizes grain formation. Deeper penetration of knives stimulates new growth by cutting through creeping bentgrass stolons and Kentucky bluegrass rhizomes while removing accumulated thatch. A more aggressive, deep vertical mowing can remove a greater amount of thatch compared with other cultivation practices. Even though this is beneficial, deeper vertical mowing should not be used when the grass is growing slowly because aggressive growth is needed to fill in disturbed areas.



Figure 20. Vertical mowing.

7.3.2 Spiking/Slicing

Spiking/slicing offers a temporary reduction in surface compaction and promotes water infiltration with minimal surface damage. Slicing is faster than core aeration but is less effective. Spiking can break up crusts on the soil surface, disrupt algae layers, and improve water infiltration.

7.4 Topdressing

Sand topdressing is used to improve the soil structure. Proper topdressing programs can relieve surface compaction, dilute organic matter, help in recovery from cultural practices, increase water and air infiltration, and protect the crowns. Layering can occur in the soil if proper programs aren't followed, limiting water and air infiltration. Table 10 shows application rates for light and frequent topdressing. These applications can be applied once every 2-3 weeks for even distribution in the soil layer.

When purchasing sand for topdressing, the source, budget, texture, and overall quality of the sand is important. Angular sand performs better than spherical shaped sand for topdressing. Particle size is important. It should be matched to the existing soil particle size if possible. A coarse particle size is better than fine.



Figure 21. Topdressing.

Table 10. Light and frequent topdressing rates

| Quantity (ft ³ /1,000 ft ²) | Quantity (lbs/1,000 ft ²) | Quantity (tons/acre) | Depth of Application (inches) |
|--|---------------------------------------|----------------------|-------------------------------|
| 0.50 | 50 | 1.1 | 0.006 |
| 0.75 | 75 | 1.7 | 0.009 |
| 1 | 100 | 2.2 | 0.012 |
| 2 | 200 | 4.4 | 0.24 |
| 4 | 400 | 8.8 | 0.48 |

7.5 Rolling

Rolling can help smooth the putting surface and maintain speeds at higher HOC. Even with a raised HOC, rolling can increase ball roll by 10 percent. Light-weight rollers typically have little negative impact on soil compaction unless the practice is over utilized or is used on high silt and clay soils when saturated with water. Rolling can also be used to remove dew off the playing surface, which reduces the possibility of dollar spot.

7.6 Cultural Practices Best Management Practices

Mowing

- ❖ Follow the recommended HOC for different turf species.
- ❖ Raise HOC by at least 30% in heavily shaded areas to improve turf health.
- ❖ Routinely use plant growth regulators, if needed, to improve overall turf health in shaded environments.
- ❖ Increase HOC in times of stress such as heat, drought, or prolonged cloudy weather to increase photosynthetic capacity and rooting depth of plants.
- ❖ If turf becomes too tall, it should not be mowed down to the desired height all at once. Tall grass should be mowed frequently and HOC gradually decreased until the desired HOC is achieved.
- ❖ Mowing frequency should increase during periods of rapid growth and decrease during dry, stressful periods.
- ❖ Decrease mowing frequency and/or increase HOC when the turf is stressed.
- ❖ Alter mowing patterns on all locations of the golf course.
- ❖ Alter the directions of mow on greens every time it is mowed.
- ❖ Rarely use inefficient mowing patterns (e.g. 9-3) on areas other than putting greens to save time, fuel, and labor.
- ❖ Return clippings to the canopy whenever possible.
- ❖ Use compressed air to blow off clippings from mowing equipment over grassed areas and before washing equipment.

- ❖ Collect pesticide-free clippings and compost or distribute in natural areas, away from surface waters.
- ❖ Use proper mowing equipment.
- ❖ Regularly sharpen and adjust blades.

Aeration

- ❖ Manage thatch levels to not become excessive. Annually monitor thatch accumulation. When thatch levels are becoming a concern, core aeration programs should be designed to remove 15% to 20% of the surface area.
- ❖ High traffic areas may require a minimum of two to four core aerations annually.
- ❖ Core aeration should be conducted only when grasses are actively growing to aid in quick recovery of surface density; midsummer for buffalograss and spring/fall for cool season grasses.
- ❖ Aeration events should be as deep as practical to prevent development of compacted layers in the soil profile as a result of cultivation.
- ❖ Consider timing of core aeration to avoid the period of *Poa annua* (annual bluegrass) seed head formation.
- ❖ Backfill holes with new root-zone materials if a drill-and-fill machine is used.
- ❖ High pressure water injection can be applied once every 3-4 weeks throughout the summer.

Surface Cultivation

- ❖ Initiate vertical mowing when thatch level reaches 0.25-0.5” in depth. Shallow vertical mowing should be completed at least monthly on putting greens to prevent excessive thatch accumulation.
- ❖ Aggressive or deep vertical mowing should not be used when the turf is growing slowly.
- ❖ Frequent shallow vertical mowing on putting greens prevents excessive thatch buildup and grain formation.

Topdressing

- ❖ Use light and frequent topdressing applications following aeration, according to Table 10.
- ❖ Use sand particle size distribution similar to the existing soil to avoid layering.
- ❖ Know the sand source, particle size ranges and ensure that sand is weed-free, uniform, and of appropriate quality.

Rolling

- ❖ Rolling of putting surfaces following mowing can increase putting speeds and allow for improved ball roll without lowering HOC.
- ❖ Avoid rolling on saturated soils to avoid compaction.
- ❖ Use lightweight rollers to minimize potential compaction.

8 INTEGRATED PEST MANAGEMENT

IPM is a sustainable approach to managing pests by combining biological, cultural, physical, and chemical tools in a way that minimizes economic, health, and environmental risks. IPM utilizes regular monitoring and record keeping to determine if and when treatment is needed. This approach takes the form of a combination of strategies and tactics to prevent unacceptable damage. When warranted, pesticides can be used after a considered selection process and applied following state and federal regulations.

8.1 Regulatory Considerations

As described in detail in the next chapter (“Pesticide Management”), pesticide usage needs to follow state and federal regulatory requirements.

Controlled burns may require local permits. For example, the SDDA's [Wildland Fire Division](#) Suppression District requires a [permit for open burns](#) conducted in the Black Hills Forest Fire Protection District.

8.2 IPM Overview

IPM is comprised of a range of pest control methods or tactics designed to prevent pests (insects, pathogens, nematodes, weeds, etc.) from reaching economically or aesthetically damaging levels while creating the least risk to the environment. IPM programs have basic components that provide the opportunity to make informed decisions on the control of pests at a golf course. Five steps for an effective IPM program for turf are as follows:

Step 1: Monitor pests and their damage and record information.

Step 2: Identify pests and understand their biology.

Step 3: Determine action threshold levels.

Step 4: Consider a variety of control methods (biological, chemical, cultural, manual, and mechanical).

Step 5: Evaluate the IPM program.

IPM is flexible, and superintendents can usually balance course quality and environmental goals through its implementation. Growing healthy turf is the best and first line of defense against pests. For example, cultural conditions that predispose turfgrass to diseases include close mowing, inadequate or excessive nitrogen fertility, frequent or excess irrigation, inadequate thatch management, poor drainage, and shade. Following cultural BMPs and nutrient BMPs can help alleviate these conditions. However, under the right conditions, pests can sometimes cause excessive damage to highly managed turfgrass.

A number of non-chemical and chemical control options are available. When plant protectant chemicals are needed, selection of an appropriate pesticide should follow an evaluation process

that considers potential impacts on beneficial organisms and the environment, as well as the potential for development of pesticide resistance. Pesticide products should be rotated, based on their resistance classification, as discussed in further detail in section 8.6.5 of this chapter.

Additional information and resources on IPM can be found on the [North Central IPM Center](#) website.

8.3 Monitoring Pests and Recording Information

In the IPM plan, pest monitoring or “scouting” efforts should be described for all areas of the course such as putting greens, tees and fairways, roughs, and landscaped areas. Scouting methods include visual inspection, soil sampling, soap flushes, and trapping for insects. Additional monitoring efforts can include weather tracking, which is especially helpful for predicting potential disease outbreaks. Here is one potential scouting schedule: daily on putting greens, at least weekly on tees and fairways, twice a month on roughs, and whenever the potential for pests increases due to weather. For example, warmer temperatures combined with high humidity favor the development of diseases such as dollar spot and brown patch.

When pests are discovered during monitoring, the pest pressure and thresholds for action should be quantified with measurements such as:

- Number of insects per unit area.
- Disease patch sizes.
- Percent of area affected.

Documentation should include useful information, such as photographs, delineation of pest boundaries on an area map, outbreak date, description of the prevailing weather conditions, and recent management practices. This information can be used to build a database for reference in future seasons and for updating the IPM plan.

8.4 Identifying and Understanding Pests

Once detected, pests must be properly identified. Understanding the biology of pest species and their vulnerable life stages assists in later control efforts. Just as important as identifying pests is recognizing and understanding beneficial organisms and their life cycles so their populations are not unduly negatively affected while managing pests. Superintendents and staff should continually hone their diagnostic skills by attending training seminars and field days, obtaining reference materials, and providing peer-to-peer training.

The following are resources for assistance in identifying and understanding pest species:

- [Turfgrass Weed Control for Professionals](#), which includes color images and descriptions of many turf weeds.
- [Managing Turf and Landscape Weeds](#).
- [Turf iNfo webpage](#).

8.4.1 Diseases

In many cases, diseases develop when conditions are favorable, regardless of management strategies. However, the severity of disease is often greatly reduced by using cultural, biological, and genetic techniques. As a rule, healthy, well-managed turf better withstands disease outbreaks and recovers more rapidly than unhealthy turf. Some common diseases in South Dakota include brown patch, [dollar spot](#), summer patch, powdery mildew, snow molds, [pythium](#) take-all patch, fairy ring, and anthracnose. If unrecognized diseases occur, samples should be sent to the [SDSU Plant Diagnostic Clinic or other contracted diagnostic labs](#).

8.4.2 Weeds

High-quality turfgrass outcompetes seedling weeds for light, water, and nutrients, and thus prevents them from establishing large weed stands that decrease turf playability and aesthetics. Weeds can also harm turf by hosting other pests such as plant pathogens, nematodes, and insects.

The potential for invasive weeds can be limited through implementation of the BMPs identified in this document related to turfgrass selection, nutrient management programs, irrigation, and cultural practices. Some common South Dakota turfgrass weeds include crabgrass, yellow nutsedge, annual bluegrass, and broadleaf weeds. The [Turfgrass Weed Control for Professionals](#) guide provides detailed weed identification and management information.



Figure 22. Dollar spot.



Figure 23. Snow mold.



Figure 24. Summer patch.

8.4.3 Nematodes

Plant-parasitic nematodes adversely affect turfgrass health by debilitating the root system of susceptible species, thus decreasing the efficiency of water and nutrient uptake. Turf weakened by nematode infestations favors further pest infestation, especially weeds. Over time, turf in the affected areas thins out and, with severe infestations, may die. Turfgrass often begins showing signs of nematode injury when stressed (e.g. drought, high or low temperatures, and wear).

8.4.4 Insects/White Grubs

Insects such as billbugs, chinch bugs, mealy bugs, webworms, armyworms, cutworms, and ants impact turfgrass in South Dakota. Armyworm and cutworm larvae feed at night on many varieties of turfgrass on the surface and then rest during the day. They are easier to detect using visual inspection and other methods based on irritating detergent-based solutions (soap flushes) to assess larval numbers.

White grubs are the larval stage of a group of beetles collectively known as scarabs (family Scarabaeidae). Among the white grub species causing turf injury in South Dakota are:

- May/June beetles, *Phyllophaga* spp. (three-year grubs).
- Northern masked chafers, *Cyclocephala* spp. (annual grubs).
- Black turfgrass ataenius, *Ataenius spretulus*.
- Japanese beetle, *Popillia japonica*.

White grubs can destroy significant areas of turfgrass, with damage appearing in the later part of summer (August through early September). Summer drought stress and insufficient irrigation may compound the damage to turf by grubs. Additional information on their biology and control methods is available on UNL's [White Grub Management](#) webpages.



Figure 25. Damage to turf caused by animals digging for grubs.

8.5 Determining Threshold Levels

A key feature of IPM programs is the identification of tolerance thresholds. Thresholds are based on the pest population, the stage of the pest, and the life stage of the plant. Injury thresholds represent the pest population level that causes unacceptable injury. Treatment thresholds are less than the injury threshold and indicate the number of pests or level of damage that would justify treatment to prevent the pest population from progressing and causing unacceptable turf loss.

8.6 Control Methods

Once a pest problem reaches the established treatment threshold, different methods can be used to control the problem, including cultural, mechanical, biological, and chemical. Selecting the most appropriate approach depends on a number of factors, including the site-specific location on the golf course, efficacy of non-chemical controls for the particular situation, economics, and pest populations.

8.6.1 Cultural Controls

Cultural practices, especially irrigation, mowing, topdressing, core cultivation, and venting, greatly affect both short- and long-term plant health. Using and/or altering cultural practices, especially in times of stress, to keep plants and soil healthy will help turf to better withstand pest pressure. It is important to recognize that turfgrass management practices such as core aeration and sand topdressing, while beneficial, can also stress turfgrass.

As an example of the role of non-chemical controls in disease management, the following practices can reduce the incidence and severity of dollar spot:

- Planting resistant cultivars. Consult [National Turfgrass Evaluation Program \(NTEP\)](#) trial results to evaluate dollar spot resistance.
- Minimizing moisture stress and leaf wetness.
- Removing morning dew as early as possible.
- Rolling putting greens three or more times per week.
- Applying biological organisms known to suppress dollar spot, such as *Bacillus licheniformis*.

8.6.2 Mechanical or Physical Controls

Mechanical methods, such as vacuuming, or physical control methods, such as hand pulling weeds, exclude or remove pests, though these methods may be time consuming and work best when pest populations are low.

8.6.3 Controlled Burns

As many golf courses convert maintained turfgrass areas to native grassed sanctuaries, many facilities use prescribed or controlled burns to reduce undesirable plants and encourage desirable species, enrich wildlife, and remove excessive plant debris. Prescribed burning is especially effective in suppressing cool-season grasses and woody plant materials to create a more desirable stand of a links-style course that resembles a tallgrass prairie. Use of a prescribed burn, along with other control methods, is an IPM approach to effectively manage these eco-sensitive areas.



Figure 26. Controlled burning at Elkhorn Ridge Golf Club.



Figure 27. Controlled burns are used to manage native grass areas.

8.6.4 Biological Controls

The biological component of IPM involves the release and/or conservation of natural predators, such as parasites and pathogens, and other beneficial organisms. Several organisms known to have some efficacy against turfgrass pests have been marketed as pest control products, such as such as *Bacillus licheniformis*. Natural enemies (e.g. ladybird beetles, green lacewings, and mantids) of some insect pests may be purchased and released near pest infestations. Areas on the golf course can also be modified to better support natural predators and beneficial organisms, especially in landscaped areas.

8.6.5 Pesticides/Chemical Controls

Chemical control is an acceptable IPM practice when other methods will not alleviate the pest problem. Reduced-risk pesticides and biopesticides provide a number of advantages over conventional pesticides and should be considered if applicable. The selection of conventional pesticides should follow a selection process and these criteria:

- They must be registered for use in South Dakota. The SDDA maintains a database of registered pesticide products. This database is searchable via [company](#) or [product](#).
- They should be effective in treating the pest problem.
- Pesticide rotation, based on resistance classification, as classified by [Fungicide Resistance Action Committee \(FRAC\)](#), [Herbicide Resistance Action Committee \(HRAC\)](#), and [Insecticide Resistance Action Committee \(IRAC\)](#).
- Costs should be considered.
- Environmental risk and potential for water quality impacts must be evaluated.
- Evaluating the environmental risk and potential for impact on water quality can include the use of software, such as the [Windows Pesticide Screening Tool \(WIN-PST\)](#), which was developed by the U.S. Department of Agriculture's Natural Resources Conservation Service to evaluate the potential of pesticides to move with water and eroded soil/organic matter and to affect non-targeted organisms. WIN-PST users can select combinations of active ingredient, soil type, and growing conditions to select an active ingredient that has less potential to leach and/or runoff into surface water.
- The use of all pesticides should follow the label and adhere to state and federal regulations, as described in the “Pesticide Management” chapter.

For more information, see:

- [UNL publications](#) for specific pest control methods.
- [UNL's Turf iNfo page](#) for recommendations on dealing with various pests.

Reduced Risk Pesticides and Biopesticides

The EPA's [Conventional Reduced Risk Pesticide Program](#) registers reduced-risk pesticides, which are commercially viable alternatives to conventional pesticides. The SDDA maintains a database of registered pesticide products that includes reduced risk pesticides.

The EPA characterizes the advantages of reduced-risk pesticides as follows:

- Low impact on human health.
- Lower toxicity to non-target organisms (birds, fish, and plants).
- Low potential for groundwater contamination.
- Low use rates.
- Low pest-resistance potential.
- Compatibility with IPM practices.

Biopesticides, which are derived from such natural materials as animals, plants, bacteria, and certain minerals, are classified separately by the EPA. For more information on biopesticides, see the EPA's [Biopesticide Registration](#) page.

8.7 Evaluation and Record Keeping

It is essential to record the results of IPM-related efforts to develop historical information, document patterns of pest activity, and evaluate successes and failures. Records of pesticide use are required by South Dakota law for restricted-use pesticides. For IPM purposes, records should be kept for all pesticide applications and should include additional information, such as monitoring records, weather records, cultural management logs, and pest response.

8.8 IPM Best Management Practices

- ❖ Develop a facility-specific, written IPM plan. Available resources for writing an IPM plan include the Golf Course Superintendents Association of America's IPM information and online tools.
- ❖ Select turfgrass cultivars and species recommended for use in South Dakota and best suited for the intended use and environmental conditions of the specific site.
- ❖ Correct the soil's physical and chemical properties that may impact turfgrass health and its ability to resist pests.
- ❖ Evaluate the potential impact of the timing of cultural practices and nutrient applications on the incidence of pest problems.
- ❖ Use a defined pesticide selection process to select the most effective pesticide with the lowest toxicity and least potential for off-target movement.
- ❖ Document all IPM-related activities, including non-chemical control methods and pesticide usage.

Monitoring Pests and Recording Information

- ❖ Monitor prevailing environmental conditions for their potential impact on pest problems.
- ❖ Train personnel how to regularly monitor pests by scouting or trapping.
- ❖ Identify alternative hosts and overwintering sites for key pests.
- ❖ Assess pest damage when it occurs, noting particular problem areas, such as the edges of fairways, shady areas, or poorly drained areas.
- ❖ Document when the damage occurred. Note the time of day, date, and flowering stages of nearby plants.
- ❖ Map pest outbreak locations to identify patterns and susceptible areas for future target applications.

Identifying and Understanding Pests

- ❖ Identify key pests in the IPM plan.
- ❖ Determine the pest's life cycle and know which life stage to target (e.g. for insect pests, whether it is an egg, larva/nymph, pupa, or adult).
- ❖ Identify pests accurately. For diseases, correctly identifying the disease pathogen often involves sending samples to diagnostic labs at land grant universities (e.g. [SDSU's plant diagnostic clinic](#), [UNL's diagnostic clinic](#)).

Determining Thresholds

- ❖ Establish injury and treatment thresholds levels for key pests and document them in the IPM plan.

Control Methods

- ❖ Implement proper cultural, irrigation, and turf management practices to reduce turfgrass stress and conditions that favor pest development.
- ❖ Maintain a proper fertilization schedule to improve turf density and quality and reduce pest populations.
- ❖ Make sure your materials, such as topdressing, are pest-free.
- ❖ Apply a preventative pesticide to susceptible turfgrass when unacceptable levels of disease are likely to occur.
- ❖ Address damage from turfgrass pests such as diseases, insects, nematodes, and animals to prevent density/canopy loss to broadleaf weeds.
- ❖ Divert traffic away from areas that are stressed by insects, nematodes, diseases, or weeds.
- ❖ When nematode activity is suspected, an assay of soil and turfgrass roots is recommended to determine the extent of the problem.
- ❖ Consider using insect-parasitic nematodes to naturally suppress insect pests such as white grubs.
- ❖ Identify areas on the golf course that can be modified to attract natural predators, provide habitat for them, and protect them from pesticide applications.
- ❖ Install flowering plants that can provide parasitoids with nectar or sucking insects (aphids, mealybugs, and soft scales) with a honeydew source.

- ❖ Avoid applying pesticides to roughs, driving ranges, or other low-use areas to provide a refuge for beneficial organisms.
- ❖ Follow a selection process when conventional pesticide use is warranted.
- ❖ Follow guidelines and advice provided by [FRAC](#), [HRAC](#), and [IRAC](#).
- ❖ Evaluate use of reduced-risk pesticides and biopesticides to treat the problem.

Controlled Burns

- ❖ Follow local permitting requirements.
- ❖ Notify the following of when/where an open burn will occur: local fire department, municipality nearest the burn, the county sheriff's department and any military, commercial, county, municipal or private airport or landing strip that may be affected by the open burn.
- ❖ Make sure the burn area is free of any debris.
- ❖ Ensure that the prevailing winds during the burn are away from any town/city or any occupied residence likely to be affected by the smoke to the best extent possible.
- ❖ Minimize the amount of dirt in the material being burned to reduce smoldering.
- ❖ Conduct open burns between three hours after sunrise and three hours before sunset to allow for smoke dispersion and to avoid air inversions that can trap the smoke at breathing level. Additionally, fuel should not be added outside the timelines listed above.
- ❖ Extinguish an open burn completely to ensure smoldering of material does not persist.
- ❖ Do not create a traffic hazard on any public road or airport right of way or obscure visibility.
- ❖ Use common sense precautions, such as having someone watching the fire until it is extinguished and assuring smoke does not impact residences or impair vehicular travel on highways.

Evaluation and Record Keeping

- ❖ After treatment, determine whether the corrective actions reduced or prevented pest populations, were economical, and minimized risks. Record and use this information when making similar decisions in the future.
- ❖ Observe and document turf conditions regularly, noting which pests are present, so that informed decisions can be made regarding the damage the pests are causing and what control strategies are necessary.

9 PESTICIDE MANAGEMENT

Pesticides are plant protectant chemicals and their use should be part of an overall pest management strategy that includes biological controls, cultural methods, pest monitoring, and other applicable practices. When a pesticide application is deemed necessary, its selection should be based on effectiveness, toxicity to non-target species, cost, site characteristics, and its solubility and persistence in the environment.

Storage and handling of pesticides in their concentrated form poses the highest potential risk to groundwater and surface water. For this reason, it is essential that facilities for storing and handling pesticides be properly sited, designed, constructed, and operated in accordance with federal and state regulations.

9.1 Regulatory Considerations

The Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) is the federal law regulating the manufacture, distribution, sale, and use of pesticides.

The EPA has given the SDDA the authority to regulate pesticides in South Dakota under FIFRA on behalf of the EPA. The SDDA is also responsible for the state's pesticide law ([SDCL 38-20A](#)), agricultural pesticide application law ([SDCL 38-21](#)), and pesticide administrative rules ([ARSD 12:56](#)). State regulatory requirements include the following:

- Bulk pesticide storage facilities require a [permit](#) from the SDDA, and storage requirements apply.
- Each pesticide operational area using containment must be registered with the SDDA. An “Operational Area” is any place where:
 - Pesticides are transferred between containers.
 - Including transfer to application equipment.
 - Pesticides are loaded, unloaded, mixed, repackaged, or refilled.
 - Pesticides are cleaned, washed, or rinsed from application, storage, handling, or transportation equipment.
 - Areas where pesticides are received or distributed in the manufacturer’s original containers that remain sealed are not considered to be operational areas.

The [Operational Area Containment Registration Application](#) must be completed and submitted to the SDDA. More information on operational area containment on the SDDA’s [Pesticide Operational Area and Containment FAQs](#) web page.

- Both commercial and private applicators who conduct operational area activities must use a written pesticide handling and discharge response plan. The plan must be kept current at all times and available for employees and inspection by the SDDA at either the operational area, the applicator's nearest local office, or the location from which the operational area is administered.

- Only commercial pesticide applicators to maintain pesticide application records, but both commercial and private applicators are required to keep records of restricted-use pesticides applications for two years.
- Pesticide must be handled, stored, and disposed of in accordance to [ARSD 12:56:15](#) and [ARSD 12:56:02](#).

The SDDA is also the lead agency responsible for administering the pesticide certification program. It trains, certifies, and licenses pesticide applicators and licenses pesticide dealers. [Commercial applicator](#) certification and licensure is valid for two years and must be renewed by March 1. The SDDA hosts annual applicator certification classes and recommends the [National Pesticide Applicator Certification Core Manual](#) as a resource for applicators seeking certification.

9.2 Human Health Risks

Pesticides belong to numerous chemical classes that vary greatly in their toxicity. Acute toxicity refers to a single exposure by mouth, skin, or inhalation, or repeated exposures over a short time. Chronic toxicity effects are associated with long-term exposure to lower levels of a toxic substance, such as ingestion in drinking water. Pesticide toxicity and level of exposure can be a risk to human health. This idea is expressed by the risk formula: Risk = Toxicity x Exposure. Therefore, risk can be held to an acceptably low level if the amount of exposure is kept low. A number of measures can be taken to mitigate risk of exposure, including:

- Reading the pesticide product label and complying with all directions.
- Dressing appropriately and using personal protective equipment (PPE).
- Storing, handling, mixing, and applying pesticides with caution and following all safety precautions.

Pesticide labels provide information on PPE and first-aid information specific to the product. Therefore, applicators should always read and follow the label before using a pesticide, in addition to following standard safe practices. Safety Data Sheets (SDS; formerly called Material Safety Data Sheets [MSDS]) provide important information on hazardous chemicals. In addition, exposure to pesticides can be mitigated by practicing good work habits and adopting modern pesticide mix/load equipment (e.g. closed loading) that reduce potential exposure.

Potential routes of exposure to golfers include exposure via shoes, clothing, and equipment. Pesticide labeling addresses re-entry restrictions, and any application should be allowed to thoroughly dry before play resumes.

For more information on human health related pesticide issues, see [Acute Pesticide Poisoning](#) on the South Dakota Department of Health website and the UNL Extension publication [Managing the Risk of Pesticide Poisoning and Understanding the Signs and Symptoms of Pesticide Poisoning](#).

9.3 Personal Protective Equipment

PPE protects workers from exposure through one or more pathways: skin, eyes, oral ingestion, or respiratory tract. Pesticide labels list legal requirements for minimum PPE, such as specific types of clothing, goggles, and respirators. The type of PPE needed depends both on the toxicity of the pesticide and the formulation. If a pesticide label does not have specific PPE requirements, the route of entry and other information on the label can be used to determine the type and degree of appropriate protection. To avoid contamination, PPE should not be stored in a pesticide storage area. For more information on PPE, see the following UNL Extension publications:

- [*Protective Clothing and Equipment for Pesticide Applicators.*](#)
- [*Pesticide Safety: Choosing the Right Gloves.*](#)
- [*Respirators for Handling Pesticides.*](#)

9.4 Environmental Fate and Transport

Environmental characteristics of a pesticide can often be determined by the environmental hazards statement found on pesticide product labels. The environmental hazards statement (referred to as “Environmental Hazards” on the label and found under the general heading “Precautionary Statements”) advises the user of product specific concerns. Potential environmental impacts include toxicity to non-target organisms and contamination of surface water or groundwater. If endangered species are present on or near the course, labeling on applicable pesticide products directs users to the limitations found in the EPA's [Endangered Species Protection Bulletins](#).

The key to preventing pesticide impacts to water quality is an understanding of the physical and chemical characteristics that determine a pesticide's interaction with the environment: solubility, adsorption, persistence, and volatilization. Pesticide characteristics influence the potential for runoff, leaching, or drift. Once applied, pesticides can move off-site in several ways: in water, in air, attached to soil particles, and on or in objects, plants, or animals.

To prevent pesticides from moving off-site, pesticide characteristics, site-specific characteristics, and prevailing conditions should all be evaluated. Pesticide characteristics, such as solubility, and site-specific characteristics, such as soil type, depth to the water table, geology, and proximity to surface water, should be considered before selecting and applying pesticides. Prevailing weather conditions, such as the chance of precipitation, the prevailing wind, and humidity, should be evaluated with respect to the timing of pesticide applications.

9.4.1 Leaching and Runoff

Most pesticide movement in water is either by surface movement off the treated site (runoff) or by downward movement through the soil (leaching). Runoff and leaching may occur when:

- Too much pesticide is applied or spilled onto a surface.

- Too much rainwater or irrigation water moves pesticide through the soil off-site or into groundwater.
- Highly water-soluble or persistent pesticides are used.

Pesticide movement in soil and water is affected by its water solubility, adsorption by soil, and persistence. Pesticides with greater adsorption by soil are less likely to be moved by leaching or surface runoff but can be carried to surface water with eroding soil. In addition to following the pesticide BMPs to reduce the likelihood of pesticides moving off-site in surface runoff, buffer strips (as discussed in the “Water Management” chapter of this document) slow down runoff and allow pesticides to adhere to soil particles and plant tissue, preventing contamination of surface water.

Pesticides with less adsorption by soil are more likely to leach through the soil and reach groundwater. For example, if rainfall is high and soils are permeable, water that carries dissolved pesticides may take only a few days to percolate down to the groundwater.

9.4.2 Drift

Air movement causing pesticide transfer away from the application site is called drift. Pesticides may be carried off-site in the air as spray droplets, vapors, or even on blowing soil particles.

- Spray drift: Airborne movement of pesticide particles to non-target sites during application.
- Vapor drift: Volatilization of particles from plant and other surfaces during and after application and movement as a gas or vapor to a non-target site in sufficient concentrations to affect plant processes.

The potential for spray drift is strongly related to droplet size; smaller droplets have smaller mass and remain airborne and exposed to air movement longer than larger droplets. Equipment selection and operation characteristics, such as nozzle type, spray pressure, nozzle spray angle, and spray volume, impact the potential for spray drift. Weather related considerations that can influence the potential for spray drift include wind speed, wind direction, air stability, relative humidity, and temperature.

The formulation of combination products as an amine or ester can also impact the potential for drift. Esters have higher vapor pressures than amines, but typically provide better weed control. In cooler weather, ester formulations can often be used safely. In higher temperatures, the risk of volatilization increases and calls for switching to an amine formulation if drift is a concern.

Vapor drift can sometimes be difficult to predict and depends on the factors such as the pesticide’s chemical characteristics and weather, even days after the application. Volatility increases as the pesticide’s vapor pressure increases and as air temperature and wind speed increase. Irrigating shortly after surface application of volatile pesticides reduces the potential for vapor drift.

To avoid either kind of drift, wind speeds of 3-10 mph are best for applying pesticides. Less than 3 mph indicates stagnant air and the potential for temperature inversions, while more than 10 mph indicates an increasing potential for particle drift. Temperature inversions can result in long distance drift. They occur when lighter warm air rises upward into the atmosphere and heavier cooler air settles near the ground. Under these conditions, air does not mix, and spray droplets do not disperse. Any subtle airflow can move this mass of pesticide spray droplets off-target. Temperature inversions typically start at dusk and break up around sunrise as air mixes vertically.

Drift management directions are typically an integral component of product labeling. Therefore, the pesticide label should be reviewed for specific information on drift reduction techniques or requirements. Weather-related instructions on the label must be followed as well.

For more information on preventing drift, see the UNL Extension publication [Spray Drift of Pesticides](#). Some specialty crops are especially sensitive to pesticides. Therefore, pesticide applicators can check the [DriftWatch](#) and [BeeCheck](#) websites, which are online mapping services from [FieldWatch](#) that allow those with commercial specialty crops, organic crops, beehives, and other sensitive crops to report their field locations. All applicators applying pesticides outdoors are encouraged to sign up for free access to the FieldCheck app and/or free email notices.

9.5 Pesticide Record Keeping

Maintaining accurate records of pesticide-related activities (e.g. purchasing, storage, inventory, and applications) is essential. South Dakota requires only commercial pesticide applicators to maintain pesticide application records. Both commercial and private applicators are required to keep records of restricted-use pesticides applications for two years. An [example pesticide application record form](#) is published by SDDA. Though not required for general use pesticides, keeping records of pesticide applications is a good practice for turfgrass management. More information on pesticide record keeping can be found on the SDDA [Pesticide Recordkeeping Requirements](#) webpage.

9.6 Pesticide Storage and Handling

Storage and handling of pesticides in their concentrated form poses the highest potential risk to groundwater or surface waters. For this reason, it is essential that care be taken in transporting pesticides and that the facilities for storing and handling these products be properly sited, designed, constructed, and operated. Guidelines for storing and handling pesticides are as follows:

- Avoid the problem of storing pesticides by purchasing only the amount needed for the current season.
- Store pesticides in their original container with the original label attached. Read each label to determine suitable storing conditions.

- Do not store pesticides with food, feed, seed, planting stock, fertilizers, veterinary supplies or pesticide safety equipment. Also, avoid storing them next to a water supply.
- Date containers as they are purchased and keep an inventory list so outdated material can be disposed of.
- Designate a building, room, or cabinet specifically for pesticide storage and nothing else. The optimum storage facility should have a concrete floor, which is impermeable and easy to wash; adequate ventilation to avoid extreme heat and to reduce the concentration of toxic or flammable vapors; insulation and supplemental heating if required to meet label specifications; good lighting; and access to water to handle accidental spills.
- Always keep the building, room or cabinet where pesticides are stored locked when the area is unattended.
- Post caution signs at all entrances or doors that warn the area is used for pesticide storage.
- Routinely examine pesticide containers for leaks, corrosion, breaks, and tears. Clean up spills immediately and properly dispose of containers and cleaning materials. Sawdust, industrial absorbent, cat litter, or dry soil may be used to soak up liquid spills. Sweeping compound can be used with dry spills. Keep cleaning materials in the storage area for quick access.

As discussed in the regulatory section, a [permit](#) from SDDA is required for a bulk pesticide storage facility. Permanently sited bulk pesticide storage containers that are able to hold more than 300 gallons (liquid) or 100 lbs (dry) are required to be placed within a bulk pesticide storage facility that includes secondary containment. A containment area for a bulk pesticide storage area must be capable of holding 110 percent of the largest volume tank plus the area displaced by the butts of all other tanks inside the containment. For more information on storage requirements, see the SDDA publication [Fertilizer and Pesticide Bulk Facility Manual](#).

A chemical handling and discharge response plan is also required. The plan needs to be specific to each facility, and a current, written plan must be available: 1) for inspection by the department, 2) for use at the operational area, 3) at the applicator's nearest local office/residence, or 4) at the location where the operational area is administered. The plan must contain the following information:

- Methods and procedures for the transfer, loading, unloading, mixing, repackaging, and refilling of pesticide containers and pesticide application equipment.
- Methods and procedures for the periodic inspection of equipment used to transfer or hold pesticides and for the repair of any equipment found to be defective.
- Methods and procedures for the rinsing, washing, and cleaning of pesticide containers and application, storage, or transportation equipment.
- Methods and procedures used in the transfer, handling, storage, and disposal of materials recovered from within operational area containment, if required.

- Methods, procedures, materials, and equipment used to contain, recover, store, transport, and dispose of discharges outside of operational area containment systems.
- The identity and telephone numbers of responsible persons and agencies to be contacted if a discharge occurs.

For more information, see the SDDA's [Private Applicator Handling and Discharge Response Plan](#).



Figure 28. Chemical storage area at Hillsvie Golf Club.



Figure 29. Pesticide mixing and loading using the appropriate PPE.

9.7 Pesticide Application Equipment and Calibration

Application equipment must apply the pesticide to the intended target at the proper rate. Information on the pesticide label specifies the legal application rate and sometimes suggests the appropriate equipment for use with the product.

To apply liquid or granular pesticides at the proper rate, properly calibrated application equipment is essential. Such equipment mitigates environmental and human health concerns, reduces the chances of over-or under-applying pesticides, and optimizes pesticide efficacy. In addition, applicators must be especially careful to avoid exposure through inhalation when applying granular products. Equipment should also be checked frequently for leaks and malfunctions.

For spray applications, the size of the equipment (tank size, boom width, etc.) should be matched to the scale of the facility. Nozzle selection and coverage, in particular, are important in the control of drift. The type of nozzle, nozzle orifice size, sprayer pressure, and the height or distance of the nozzles from the target affect the potential for off-site movement of pesticides. A nozzle that primarily produces coarse droplets is usually selected to minimize off-target drift.

For more information on selecting application equipment to prevent drift, see the UNL Extension publication [*Spray Drift of Pesticides*](#).

9.8 Transportation

According to state law, it is a violation to transport pesticides in any manner that will endanger humans, animals, or the environment. SDDA recommendations for pesticide transportation are as follows:

- Use a ratchet-type tie down strap or chain binder to secure tanks to the vehicle. Make sure that the strap or chain/chain binder is of sufficient strength to secure the load.
- Inspect all plumbing and secure hoses and other equipment to avoid damage and potential spills.
- Insure that the transport vehicle is capable of transporting the weight of the container and contents.
- When transporting small containers:
 - Do not transport them inside the passenger compartment.
 - It is suggested that small pesticide containers be placed within a leak proof container such as a covered plastic container.
 - Never leave pesticides unattended in an unlocked vehicle or an unsecured area where they can be tampered with or stolen.
 - Secure pesticide containers in an area of the vehicle to avoid significant movement or breakage from movement of other items in the vehicle.

Spills during transport are required to be reported to the Department of Agriculture or Emergency Management Services within 12 hours after a spill of more than 5 gallons of liquid or 50 pounds of dry pesticide.

9.9 Mixing/Washing Station

Procedures for mixing of pesticides and washing of pesticide application equipment will be detailed in the written pesticide handling and discharge response plan. For detailed information on cleaning application equipment, the UNL Extension publication [Cleaning Pesticide Application Equipment](#) provides detailed information on cleaning equipment and the University of Missouri Extension publication [Cleaning Fields Sprayers to Avoid Crop Injury](#) describes a sprayer cleanout procedure in detail.

9.10 Disposal

The safest way to dispose of leftover pesticide from professional applications is to use all of the chemical according to directions on the label. This includes the washwater from pesticide equipment washing, which must be used in accordance with the label instructions.

Often pesticide storage facilities accumulate unusable or unwanted pesticide products. They can accumulate for a variety of reasons, such as mistakes made in calculating the amount of product needed or the launch of new product chemistries that may be more effective at controlling target pests. Disposing of these stockpiles properly may be challenging. Simply keeping them in storage eventually becomes problematic when packaging inevitably deteriorates or corrodes and creates a hazard.

In 1992, the South Dakota Legislature enacted laws allowing the creation of programs for pesticide container recycling and unusable-pesticide collection projects. SDDA administers the programs, with cooperation from the SDSU Extension, the agriculture industry, local entities, and individuals. The SDDA [Container Recycling & Waste Pesticide Collection](#) webpage provides information, collection dates, and registration forms. The SDSU Cooperative Extension publication [Waste Pesticides: Proper Storage, Handling and Disposal](#) offers more information on proper procedures.

9.11 Pesticide Container Management

Handling of empty pesticide containers must be done in accordance with label directions as well as with all laws and regulations. Under the federal Resource Conservation and Recovery Act, a pesticide container is not empty until it has been properly rinsed. Non-refillable pesticide containers that have been properly rinsed can be handled and disposed of as non-hazardous solid waste; some plastic containers can be recycled. Refillable containers may be returned to the supplier unrinsed.

Detailed procedures for cleaning pesticide containers, such as pressure rinsing and triple rinsing, and storing empty containers are described in the SDDA's [Reducing Pesticide Waste in South Dakota](#).

Pesticide containers can be either recycled through participation in the SDDA container recycling program or disposed of by depositing them in a licensed sanitary landfill after pressure rinsing or triple rinsing. Ways to reduce the amount of waste that requires handling include identifying and implementing waste-reduction practices and purchasing bulk packaging when possible.

9.12 Emergency Preparedness and Spill Response

Pesticide leaks or spills, if contained, will not percolate down through the soil into groundwater or be carried away in runoff. However, if not contained, controlled, and cleaned up properly, pesticide spills can harm human health, the environment, or both through leaching or runoff.

The required chemical handling and discharge response plan must include information on spill response and be available: 1) for inspection by SDDA, 2) for use at the operational area, 3) at the applicator's nearest local office/residence, or 4) at the location where the operational area is administered. For more information and a plan template, see the SDDA's [Fertilizer and Pesticide Bulk Facility Manual](#). It is recommended that a current copy of the plan be provided to local fire

departments, police departments, emergency medical services, and county emergency management directors.

Minor spills of pesticide or rinsate that have occurred from the handling, loading, or cleansing of bulk containers and that accumulate in the secondary containment area must be disposed of as provided by the pesticide label. In South Dakota, the following apply:

- Pesticide discharges in excess of 25 pounds active ingredients that occur at operational areas outside of operational area containment *must* be reported to the SDDA.
- The operator or manager of a bulk pesticide storage facility shall notify the SDDA or the Division of Emergency Management **within 3 hours** after a spill of more than 25 gallons of liquid or 500 pounds of dry pesticides outside the secondary containment area.
- The commercial carrier shall notify the SDDA or Division of Emergency Management **within 12 hours** after the spill of more than 5 gallons of liquid or 50 pounds of dry pesticide which occurs during transportation. The commercial carrier shall provide **written notice** of a spill to the SDDA **within 72 hours** after the spill.

For human injuries, medical emergencies, or fires, first responders/EMT should be notified.

9.13 Pesticide Management Best Management Practices

Human Health Risks

- ❖ Follow the pesticide label for re-entry period requirements or recommendations following application.
- ❖ Allow all pesticide applications to dry thoroughly before allowing play to resume.
- ❖ Prioritize using lower risk products whenever possible.

Personal Protective Equipment

- ❖ Follow pesticide labels for appropriate PPE.
- ❖ Provide adequate PPE for all employees who work with pesticides, including equipment technicians who service pesticide application equipment.
- ❖ Ensure that PPE is sized appropriately for each person using it.
- ❖ Ensure that respirators are seal- and fit-tested properly and that the person is thoroughly trained and has no medical limitations to respirator use.
- ❖ Store PPE where it is easily accessible, but not in the pesticide storage area.
- ❖ Forbid employees who apply pesticides from wearing facility uniforms home by providing laundering facilities or a uniform service.
- ❖ Meet requirements for the [OSHA 1910.134 Respiratory Protection Program](#).

Pesticide Applications

- ❖ Consider pesticide characteristics in the chemical selection process.

- ❖ Identify any areas on the course prone to leaching losses (e.g. shallow water tables, sand-based putting greens, coarse-textured soils, etc.). Do not use highly soluble pesticides in these areas.
- ❖ Select low or non-volatile pesticides.
- ❖ If listed species or species of concern are present, specifically select pesticides that have no known effects on these species.
- ❖ Check the forecast before applying pesticides and apply when conditions are favorable, such as minimal wind velocity, temperature inversions not forecast, rain not forecast, etc.
- ❖ Follow the pesticide label to avoid drift.
- ❖ Use spray additives within label guidelines.
- ❖ Schedule the timing and amount of irrigation needed to water-in products (unless otherwise indicated on label) without over-irrigating.
- ❖ If sites adjacent to the application area are planted with susceptible plants or crops, allow a buffer area between the two, or wait until winds are blowing away from the area of concern.

Pesticide Application Equipment

- ❖ Use an appropriately sized applicator for the size of area being treated.
- ❖ Ensure the spray technician is experienced, certified, and properly trained.
- ❖ Properly calibrate all application equipment at the beginning of each season (at a minimum) or after equipment modifications.
- ❖ Check equipment daily when in use.
- ❖ Use recommended spray volumes for the targeted pest to maximize efficacy.
- ❖ Calibration of walk-behind applicators should be conducted for each person making the application to take into consideration walking speed, etc.
- ❖ Avoid high spray boom pressures; consider 45 PSI a maximum for conventional broadcast ground spraying.
- ❖ Use drift-reduction nozzles that produce larger droplets when operated at low pressures.
- ❖ Use wide-angle nozzles and low boom heights and keep boom stable.
- ❖ When possible, use lower application speeds to avoid drift.

Pesticide Record Keeping

- ❖ Use electronic or hard-copy forms and software tools to properly track pesticide inventory.
- ❖ Keep and maintain records of all pesticides; records of restricted use pesticide applications are legally required to be kept for two years.
- ❖ Use records to monitor pest control efforts and to plan future management actions.

Pesticide Storage and Handling

- ❖ Routinely undergo a “risk assessment” to identify any potential risks to the applicator or environment.
- ❖ Do not transport pesticides in the passenger section of a vehicle.
- ❖ Never leave pesticides unattended during transport.

- ❖ Maintain an inventory of all pesticides used and the SDS for each chemical.
- ❖ Avoid purchasing large quantities of pesticides that require storage for more than six months.
- ❖ Adopt the “first in-first out” principle, using the oldest products first to ensure that the product shelf life does not expire.
- ❖ Locate pesticide storage facilities away from other structures to allow fire department access.
- ❖ Store, mix, and load pesticides away from sites that directly link to surface water or groundwater (e.g. wells).
- ❖ Store pesticides in a lockable concrete or metal building separate from other buildings.
- ❖ Shelving should be made of sturdy plastic or reinforced metal.
- ❖ Metal shelving should be kept painted to avoid corrosion. Wood shelving should never be used, because it may absorb spilled pesticides.
- ❖ When storing pesticides on shelves, place liquid pesticides on lower shelves and dry formulations above them.
- ❖ Store herbicides, insecticides, and fungicides in separate areas within the storage unit.
- ❖ Storage facility floors should be impervious and sealed with a chemical-resistant paint.
- ❖ Floors should have a continuous sill to retain spilled materials and should not have drains, although a sump may be included.
- ❖ Sloped ramps should be provided at the entrance to allow the use of wheeled handcarts for moving material in and out of the storage area safely.
- ❖ Automatic exhaust fans and an emergency wash area should be provided. Light and fan switches should be located outside the building, so that both can be turned on before employees enter the building and turned off after they leave the building.
- ❖ Avoid temperature extremes inside the pesticide storage facility.

Mixing/Washing Station

- ❖ Follow secondary containment requirements as required.
- ❖ Load and mix pesticides over an impermeable surface, such as a concrete pad.
- ❖ Mix pesticides at least 150 feet downslope from any well.
- ❖ Mix materials according to label directions and in amounts that will be used for the application to avoid excess that will need disposal.
- ❖ Either use anti-backflow devices when mixing pesticides or maintain a 6” air gap between mixing container and water source.
- ❖ Pump the sump dry and then clean it at the end of each day. Liquids and sediments should also be removed from the sump and the pad whenever pesticide materials are changed to an incompatible product (i.e. one that cannot be legally applied to the same site).

Disposal

- ❖ Collect washwater (from both inside and outside the application equipment) and use it as a pesticide in accordance with the label instructions.
- ❖ The rinsate may be applied as a pesticide (preferred) or stored for use for the next compatible application.

- ❖ Annually review pesticide inventories and properly dispose of unusable and unwanted pesticides.

Pesticide Container Management

- ❖ Rinse pesticide containers immediately in order to remove the most residue.
- ❖ Rinse containers during the mixing and loading process and add rinsate water to the finished spray mix.
- ❖ Rinse emptied pesticide containers by either triple rinsing or pressure rinsing.
- ❖ Use refillable pesticide containers only for pesticides.
- ❖ Recycle non-refillable containers when possible.
- ❖ Puncture empty and rinsed pesticide containers prior to disposal and dispose of them according to the label.

Emergency Preparedness

- ❖ Keep a written pesticide handling and discharge response plan as required that outlines the procedures to control, contain, and clean up spilled materials.
- ❖ Train all employees on the emergency response plan and emergency procedures.
- ❖ Provide a copy of the written handling and discharge response plan to local authorities.
- ❖ Keep an appropriate spill containment kit in a readily available space.
- ❖ For small liquid spills, use absorbents such as cat litter or sand and apply as a topdressing in accordance with the label rates, or dispose of as a waste.
- ❖ For small solid spills, sweep up and use as intended.
- ❖ Ensure that SDS documents are present and that all employees have been properly trained on their location and contents.
- ❖ Report releases to the SDDA when required.
- ❖ For larger spills, follow guidance from the SDDA for cleanup and disposal.

10 POLLINATOR PROTECTION

Wild pollinators and managed bees, such as honeybees, contribute to South Dakota's biodiversity and support the state's agricultural economy. Most flowering plants need pollination to reproduce and grow fruit. In South Dakota, pollinators include species of native bees, honey bees, beetles, flies, moths, and butterflies. These pollinators are economically important to agriculture in the state as many cultivated crops such as alfalfa and sunflowers require pollination. Plants that promote soil conservation, such as wildflowers in woodlands and meadows, also require pollination. In addition, South Dakota has a robust beekeeping industry, with South Dakota beekeepers producing a honey crop worth over \$34.6 million from 280,000 colonies in 2016.

Pollinators around the world are facing a number of threats that impact their health, abundance, and distribution. In South Dakota, some pollinator species, such as the monarch butterfly (*Danaus plexippus*) and the western bumble bee (*Bombus occidentalis*), have been identified as at risk due to declining populations. One such threat is pesticides, which can have negative effects ranging from gene expression within individuals to colony level impacts. In particular, neonicotinoids have been a focus of research with respect to their impact on pollinators and their prevalent use in agricultural and ornamental settings. At this point, the potential connection between increased neonicotinoid use and recent pollinator decline is the subject of scientific research and debate.

Pesticide applicators need to be mindful of the impact that pesticides used on golf courses may have on pollinator species and their habitat. In addition to adhering to best management practices related to pesticide management and application, golf course managers can protect and enhance habitat on the course in a number of ways to help pollinators.

SDDA has also formulated a [South Dakota Managed Pollinator Plan](#) with voluntary guidance on protecting pollinators in the state, which is aimed at protecting honey bees, but is also relevant for protecting wild pollinators.

10.1 Regulatory Considerations

Pollinator protection language is a requirement for pesticide labels, and following the label is mandatory. Pesticide applicators must be aware of honey bee toxicity groups and be able to understand precautionary statements. In addition to following legal requirements, pesticide applicators should understand the effects of pesticides on bees and other pollinators and the routes of potential exposure. The USGA publication [Making Room for Native Pollinators](#) and the U.S. Department of Agriculture's Forest Service and Pollinator Partnership publication [Bee Basics: An Introduction to Our Native Bees](#) provide basic facts about pollinator biology that are useful to pesticide applicators. In addition, recordkeeping may be required by law to use some pesticides. Many of the IPM best management practices, such as record keeping, are valuable tools for protecting pollinators.

Some golf courses maintain beehives on site as a way to increase their environmental stewardship efforts. In South Dakota, state law requires that all apiaries, including hobbyists, must be registered with the Department of Agriculture by February 1 each year, or within 10 days after coming into possession of an apiary. For more information, see the SDDA's [Apiary Inspection](#) webpage.

10.2 Pest Management Practices

Protecting pollinators on the golf course does not preclude the use of pesticides, but instead minimizes any potential impact from these chemicals. Pesticide applicators must use appropriate tools to help manage pests while safeguarding pollinators, the environment, and humans. Using IPM best management practices is an important key to protecting pollinators because they reduce pesticide usage and minimize the potential of exposure. Superintendents can utilize IPM best management practices for turf that protect pollinators by following these simple steps:

- Identifying what is truly a pest. (For example, solitary ground-nesting bees and wasps might be alarming, but most are harmless.)
- Setting higher weed thresholds in low-use areas.
- Monitoring bee activity to avoid applying pesticides during peak activity times (i.e. applying pesticides in the early morning or evening).

When the use of pesticides is necessary, being mindful of pollinators includes selecting chemicals with low toxicity to bees, short residual toxicity, or properties repellent to bees; using caution when applying near flowering plants, including flowering weeds (mow first whenever possible); and avoiding drift. In addition, applicators are also encouraged to utilize [FieldWatch](#) or [BeeCheck](#) to locate any nearby apiaries before applying pesticides on the course. For more information on best practices to protect pollinators when using chemical control, see [Best Management Practices for Turf Care and Pollinator Conservation: Fast Facts](#) and the North Dakota State university video [Protecting Bees from Insecticide Poisoning](#).

10.3 Preserving and Enhancing Habitat on the Course

Habitat for pollinators includes foraging habitat, nesting sites, and water sources. Increasing habitat to meet pollinator needs can be accomplished simply by adding to existing plantings or through more intensive efforts to establish a larger native area. Pollinator habitat on the golf course includes existing out-of-play areas (such as buffer strips around water courses and bodies of water) and areas renovated specifically with pollinators in mind that include native plants, wildflowers, and flowering trees and shrubs, nesting sites, and water sources. Out-of-play pollinator habitats have been shown to help nearly 50 species of pollinating insects, according to the article "[Operation Pollinator for Golf Courses](#)".

To convert existing out-of-play areas to a new native area, site preparation is key and may require more than one season of effort to reduce competition from invasive or other undesirable plants prior to planting. For more information on establishing a native area, see [Making Room](#)

[For Native Pollinators](#) and [Pollinator Meadow Upper Midwest Installation Guide and Checklist](#).

For information on creating habitat specifically for monarch butterflies, a species in decline, see the [Monarchs in the Rough website](#) and [Monarch Nectar Plant Guide: Northern Plains](#). The South Dakota [Habitat Pays](#) program may be able to provide financial assistance for creating pollinator habitat.



Figure 30. Milkweed plants provide habitat for monarchs.



Figure 31. Monarch caterpillar.

Pollinator-friendly habitat contains a diversity of blooming plants of different colors and heights, with blossoms throughout the entire growing season. Native plants are best for providing the most nutritious food source for native pollinators. Though wildflowers are most often thought of as pollinator-friendly plants, grasses such as big bluestem are a major pollen source for native bees in South Dakota. In addition, many sedge species are larval hosts, hollow stem grasses provide nesting habitat, and sturdy grasses shelter insects from harsh weather. Milkweed provides habitat specifically for monarch butterflies. References for selecting plants include:

- [Selecting Plants for Pollinators: A Regional Guide for Farmers, Land Managers, and Gardeners in the Prairie Parkland](#). North American Pollinator Protection Campaign and Pollinator Partnership.
- [Pollinator Plants of the Central United States Native Milkweeds \(*Asclepias spp.*\)](#). Xerces Society.

Providing nesting sites for native species can be accomplished by taking simple steps in out-of-play areas, such as:

- Leaving exposed patches of bare soil.
- Leaving dead trees, stumps, and posts.
- Planting hollow stem grass species.
- Providing stem bundles of hollow plant stems like bamboo.
- Creating bee blocks for solitary nesters such as mason and leafcutter bees.
- Creating artificial boxes for bumble bees.

Bee boxes can be purchased or constructed following simple instructions. The UNL Extension publication [*Attracting Pollinators to Your Landscape*](#) offers instructions for constructing a simple bee box.

A clean, reliable source of water is another essential habitat consideration for pollinators. Pollinators can use natural and human-made water features such as running water, pools, ponds, and small containers of water. Water sources should have a shallow or sloping side so the pollinators can easily approach the water without drowning. In addition, irrigation management practices that preserve ground-nesting pollinators include irrigating at night and avoiding flooding any areas.

10.4 Managed Bee Hives on the Course

Hosting honey bee hives on the golf course provide bees with valuable green space, especially in urban areas, and can be a positive public-relations tool. If embarking on this effort, consider:

- Partnering with an experienced local beekeeper. Proper beekeeping is time and knowledge intensive. If not partnering with an experienced beekeeper, superintendents or other responsible staff should attend a beekeeping course.
- Ensuring enough food sources are available for both honey bees and wild pollinator species.
- Placing hives away from areas where golf course workers or golfers are active to avoid stings.
- Facing the hive exit in a direction away from in play areas of the course.
- Educating golfers via explanatory signs, newsletters, and sales of honey and other bee products.
- Calling in an experienced beekeeper if disease or parasites are suspected in order to identify and mitigate any health issues.



Figure 32. Golf courses can host managed bee hives and generate honey.

10.5 Pollinator Protection Best Management Practices

Pest Management Practices

- ❖ Before applying a pesticide, inspect the area for both harmful and beneficial insect populations, and use pesticides only when a threshold of damage has been indicated.
- ❖ Consider biological control agents, lures, baits, and pheromones as alternatives to insecticides for pest management.
- ❖ When insecticides (or other pesticides) are needed, select those with a lower impact on pollinators, such as chlorantraniliprole.
- ❖ If a granular formulation will control the pest, choose it over liquid formulations. Granular versions of pesticides are known to be less hazardous to bees.
- ❖ Restrict applications to early morning or evening when pollinators are not as active.
- ❖ Avoid applying pesticides during bloom season, and mow first to remove blooms, including those of flowering weeds such as white clover.
- ❖ Avoid application during unusually low temperatures or when dew is forecast.
- ❖ Use the latest spray technologies, such as drift-reduction nozzles to prevent off-site translocation of pesticide.

Habitat Protection and Enhancement

- ❖ Follow site preparation guidelines when renovating areas to ensure success.
- ❖ Choose south-facing sites whenever possible for establishing native areas.
- ❖ Place plants in masses (three or more) to attract pollinators.
- ❖ Select plants of different shapes, sizes, and colors and that bloom at different times of the year.
- ❖ Select native grasses that provide foraging and nesting habitat.
- ❖ Use both perennials and annuals.
- ❖ Leave stems and coarse, woody debris in native areas for pollinator nesting.
- ❖ Leave exposed patches of well-drained soil in native areas for pollinator nesting.
- ❖ Provide water sources with shallow sides for pollinators.

11 MAINTENANCE OPERATIONS

It is the objective to manage the potential environmental risks associated with golf course maintenance operations. Our industry has a need and responsibility to implement, manage, measure, and improve, pertaining to all aspects of environmental stewardship. It is imperative that hazardous materials be handled, stored, recycled, and disposed in a safe, healthy, and environmentally sound manner.

Pollution prevention includes the proper delivery, storage, handling, and disposal of all chemicals, washwater, and wastewater. For example, washwater from pesticide application equipment must be managed as a pesticide. Conversely, wastewater not contaminated with harmful chemicals can be reused or discharged to a permitted stormwater treatment system. The “Pesticide Management” chapter discusses many maintenance operations-related BMPs specifically for pesticides.

For unintended releases of any chemicals, an emergency plan, spill kit, and first-aid kit should be readily available.

11.1 Regulatory Considerations

As discussed in the “Pesticide Management” chapter of this document, the SDDA regulates pesticides transportation, storage, handling, and disposal in the state. In the case of fertilizers, while South Dakota regulates fertilizer storage and handling for commercial fertilizer distributors, these requirements do not apply to golf facilities, with the exception of reporting spills.

Owners and operators of underground storage tanks (UST) and above-ground storage tanks (AST) must register with the DENR. Petroleum storage tanks for golf facilities are regulated by DENR under the Administrative Rules for South Dakota (ARSD) 74:56:01, Underground Storage Tanks and 74:56:03, Aboveground Storage Tanks. Plans and specifications for regulated AST and UST systems must be submitted to DENR for review and approval at least 30 days before beginning construction.

Petroleum spills less than 25 gallons must be cleaned up within 24 hours. If the spill is not cleaned up within 24 hours, it must be reported. Spills and/or overfills of petroleum exceeding 25 gallons, or any spill volume that is below ground surface or impacts surface water must be reported immediately to DENR and applicable local agencies. DENR’s Ground Water Quality Program is responsible for guiding the responsible parties in their efforts to clean up their releases to state standards.

11.2 Storage and Handling of Fertilizers

Storage facilities that are well designed and well maintained protect people from exposure, reduce the potential of environmental contamination, protect chemicals from extreme temperatures and excess moisture, and, in general, reduce liability concerns and potential

environmental risks. The storage area should be secure and provide containment features. In addition, storing and maintaining equipment properly extends its useful life and reduces repairs. SDDA publishes the voluntary storage guidelines in [On Farm Fertilizer Storage](#).



Figure 33. Fertilizer storage at Hillsvie Golf Club.

11.3 Equipment Washing

Equipment washing should be conducted under controlled conditions in an appropriate contained area with minimal risk to the environment to prevent adverse washwater runoff impacts whenever possible. Equipment washing guidelines and restrictions should be established that reduce the potential for pollutants to reach stormwater runoff, surface water or groundwater.

Proper cleaning of equipment helps prevent residues from reaching surface waters, groundwater, drainage pipes, or storm sewers. The residues from washing equipment include grass clippings, soil, soaps, oil, fertilizers, and pesticides.

A primary concern when washing mowing equipment is the nitrogen and phosphorus nutrients in grass clippings. Using compressed air to blow clippings off mowers before washing can help reduce the amount of nutrients that enter drains via washwater. The best practice is to have a dedicated wash area with a catch basin to collect remaining grass clippings. Clippings can be collected, then composted or removed to a designated debris area. When formal washing areas are not available, a “dog leash” system using a short, portable hose to wash off the grass at random locations, away from surface waters, wells, or storm drains, is an option.



Figure 34. Separating clippings reduces nutrients in washwater.

For equipment with possible pesticide residue, BMPs should be followed to ensure that washwater does not become pollution source. Captured washwater can be used as a dilute pesticide per label, or it may be pumped into a rinsate storage tank for use in the next application and used as a dilute pesticide per the label.

11.4 Equipment Storage and Maintenance

Like chemical storage facilities, equipment storage and maintenance facilities should be designed to prevent the accidental discharge of chemicals, fuels, or contaminated washwater from reaching water sources. Properly storing and maintaining equipment also extends the useful life of machines and reduce repairs.

11.5 Fueling Facilities

Fueling areas should be properly sited, designed, constructed, and maintained to prevent petroleum products from being released into the environment through spills or leaks. Above-ground tanks are easier to monitor for leakage and are therefore the preferred storage method. Because of the potential for groundwater contamination from leaking USTs, leak detection monitoring is a critical aspect of UST compliance. Any leaks or spills must be contained and cleaned immediately. As discussed in the "Regulatory Considerations" section of this chapter,

releases of certain quantities must be reported to DENR. The [Petroleum Assessment and Cleanup Handbook](#) provides more information on remediation of petroleum releases.

Fueling areas should be sited on impervious surfaces, equipped with spill containment and recovery facilities, and located away from surface waters and water wells. Catch basins in fueling areas should be directed toward an oil/water separator or sump to prevent petroleum from moving outside any containment structure. Floor drains in fueling areas should be eliminated unless they drain to containment pits or storage tanks.



Figure 35. Fueling station with an AST for fuel storage.

11.6 Waste Handling

Facilities need to regularly review how they handle the disposal of unwanted, expired, or accumulated items, including chemicals, paints, pesticides, tires, batteries, used oils, solvents, paper products, plastic or glass containers, and aluminum cans. Developing recycling programs reduces waste and minimizes the quantity of waste reaching landfills. In some cases, recycling of some wastes may be required locally, and superintendents should be aware of these requirements.

All packaging from chemicals, their containers and other wastes should be properly disposed of. Pesticide-specific waste handling requirements are identified on the pesticide label and are discussed in more detail in the “Pesticide Management” chapter.

11.7 Maintenance Operations Best Management Practices

Fertilizer Storage

- ❖ Review groundwater sensitivity information before constructing any fertilizer storage facilities or handling areas.
- ❖ Storage facilities should not be located in areas with high probability of flooding.
- ❖ Locate dry fertilizer storage buildings or liquid fertilizer secondary containment over 500 feet away from a well, water supply or surface water runoff area.
- ❖ Construct storage buildings to prevent seepage or spillage of fertilizer under normal conditions.
- ❖ Unless stored in a totally enclosed building, all nonliquid fertilizer materials should be covered and stored within an appropriate secondary containment storage structure.
- ❖ Construct liquid fertilizer secondary containment capable of holding 125 percent of the volume of the largest container plus the volume of the butts of all other containers inside the liquid containment area.
- ❖ Construct dry storage for secondary containment which is of sufficient thickness and strength to withstand loading conditions.
- ❖ Design loading areas to prevent spillage onto unprotected areas and create a proper cleanup area by installing curbed containment.
- ❖ Post warning signs on chemical storage buildings, especially near entry or exit areas.
- ❖ Storage facilities should be secured and allow access only to authorized staff.
- ❖ Replace worn or faulty valves, plugs, and threaded fittings in storage containers.
- ❖ Install backflow prevention devices or use air gap separation on water supply lines used for fertilizer mixing or equipment rinsing.
- ❖ Lock valves and shutoff devices while storage containers and facilities are not in use.
- ❖ Follow hazard safety rules, worker protection laws and fire prevention rules while handling and storing fertilizer.
- ❖ Apply appropriate sealant to seams and cracks in all storage facilities and load/wash/rinse pad areas.
- ❖ Use approved containers designed for and compatible with the fertilizer being stored.
- ❖ Shelves should be made of plastic or reinforced metal. Metal shelving should be coated with paint to avoid corrosion. Wood shelving should not be used due to its ability to absorb spilled chemicals.
- ❖ Exhaust fans and an emergency wash station should be provided.
- ❖ Light and fan switches should be located on the exterior of the storage facility.
- ❖ Store liquid materials below dry materials to prevent contamination from a leak.
- ❖ Train staff and other management on how to access and use the facility's SDS database.
- ❖ Maintain accurate inventory lists.

Equipment Washing

- ❖ Brush or blow off accumulated grass clippings from equipment using compressed air before washing.
- ❖ Wash equipment on a concrete pad or asphalt pad that collects the water. After the collected material dries, collect and dispose of it properly.
- ❖ Washing areas for equipment not contaminated with pesticide residues should drain into oil/water separators before draining into sanitary sewers or holding tanks.
- ❖ Do not wash pesticide-application equipment on pads with oil/water separators. Do not wash near wells, surface water, or storm drains.
- ❖ Use spring-loaded spray nozzles to reduce water usage during washing.
- ❖ Minimize the use of detergents. Use only biodegradable, non-phosphate detergents.
- ❖ Use non-containment washwater for field irrigation.
- ❖ Do not discharge non-contaminated wastewater during or immediately after a rainstorm, since the added flow may exceed the permitted storage volume of the stormwater system.
- ❖ Do not discharge washwater to surface water, groundwater, or susceptible/leachable soils either directly or indirectly through ditches, storm drains, or canals.
- ❖ Never discharge to a sanitary sewer system without written approval from the appropriate entity.
- ❖ Never discharge to a septic tank.
- ❖ Do not wash equipment on a pesticide mixing and loading pad. This keeps grass clippings and other debris from becoming contaminated with pesticides.
- ❖ Solvents and degreasers should be used over a collection basin or pad that collects all used material.

Equipment Storage and Maintenance

- ❖ Store equipment in areas protected from rainfall. Rain can wash residues from equipment and potentially contaminate the surrounding soil or water.
- ❖ Perform equipment maintenance activities in a completely covered area with sealed impervious surfaces.
- ❖ Drains should either be sealed or connected to sanitary sewer systems with the approval of local wastewater treatment plants.
- ❖ Solvents and degreasers should be stored in locked metal cabinets away from any sources of open flame.
- ❖ Complete a chemical inventory and keep SDS of each on site. A duplicate set of SDS should be kept in locations away from the chemicals, but easily reached in an emergency.
- ❖ Use PPE when working with solvents.
- ❖ Use containers with dates and contents clearly marked when collecting used solvents and degreasers.

Fueling Facilities

- ❖ Above-ground fuel tanks are preferred as they are more easily monitored for leaks as compared with underground tanks.

- ❖ Fueling stations should be located under roofed areas with concrete pavement whenever possible.
- ❖ Fueling areas should have spill containment and recovery facilities located near the stations.
- ❖ Develop a record-keeping process to monitor and detect leakage in USTs and ASTs.
- ❖ Visually inspect any AST for leakage and structural integrity.
- ❖ Secure fuel storage facilities and allow access only to authorized and properly trained staff.
- ❖ Report spills and leaks to DENR as required.

Waste Handling

- ❖ Label containers for collecting used solvents, oils, and degreasers.
- ❖ Recycle lead-acid batteries. If not recycled, batteries are classified as hazardous waste.
- ❖ Store old batteries on impervious surfaces in areas protected from rainfall.
- ❖ Recycle used tires, paper products, plastic or glass containers, aluminum cans, and used solvents, oils, and degreasers.
- ❖ Provide a secure and specifically designated storage for the collection of recyclable waste products.
- ❖ Recycle or properly dispose of light bulbs and fluorescent tubes.

12 LANDSCAPE

The fundamental principle for the environmentally sound management of landscapes is “choose the right plant, in the right place.” Ideal landscape plants are native and adapted specifically to the soil, degree and direction of slopes, precipitation type and amounts, wind direction and speed, light patterns, and microclimate. Susceptibility to major damage by insects and other pests is another selection criterion, as are the nutrient levels of the area. By using native or adapted plants that mimic natural ecosystems, landscapes that are designed for the specific location, management capabilities and desired style can reduce overall management inputs, attract pollinators, provide multi-season interest, and enhance out-of-play areas.



Figure 36. A natural rock wall used to frame a landscaping bed adds visual interest.

12.1 Planning and Design

Planning begins with a careful assessment of existing conditions. Slopes and drainage patterns impact not only the playability of the course, but the survival of existing and proposed plants. A majority of the non-play areas on the golf course should remain in natural cover. Supplemental planting of native or adapted trees, shrubs, and herbaceous vegetation can limit soil erosion,

protect stream banks, and enhance wildlife habitat, including non-game species, birds, and pollinators, in non-play natural areas. Mimicking natural ecosystems by leaving dead trees (snags), brushy understory plants, and native grasses and forbs in these areas also reduces maintenance work, by minimizing or eliminating the need to mow or apply fertilizer or pesticide.

Higher-impact, higher-use landscape areas, such as around the clubhouse, should be designed to utilize natural drainage patterns and channel runoff away from impervious surfaces (e.g. paved areas), conserve water, and lower nutritional input requirements once mature. Installing rain gardens in locations where they catch and temporarily hold water (such as near roofs and other impervious surfaces) helps control stormwater runoff, remove contaminants before releasing water into the surrounding soil or aquifer, and conserve water by reducing supplemental irrigation needs. For more information on rain gardens, see the Natural Resources Conservation Service publication [*Rain Gardens: Capturing and Using the Rains of the Great Plains*](#).

Golf courses are excellent facilities for zoning the landscape, using designations of high-impact zones, transition zones, and perimeter zones, and matching high-use and high-impact areas to plants that need more water and likely more-intense management. Taking into consideration the lay of the land, including differences in soil and changes in sunlight levels throughout the day, planning for landscaped areas should include consideration of the water needs in each area. A zoned approach is an efficient way to plan, as follows:

- High-use and high-impact zones: Match plants that need the most water to small, highly visible areas that will be watered as needed.
- Transition zones: Choose plants that require moderate amounts of water to be applied only when they show signs of moisture stress, such as wilting.
- Perimeter zones: Use plants with minimal water requirements. Water during establishment and in periods of extreme drought.

Ideally, 10% or less of the landscaped areas should be zoned for high water use, 30% or less of the area should be zoned for moderate water use, and 60% or more of the landscape should be zoned for low water use.

12.2 Site Inventory and Assessment

Before developing a landscape plan, an inventory should be conducted of existing plants, their condition and quality, their contribution to the overall style of the course, and how they've been managed. For landscaped areas, a soils analysis and a soil test should be conducted. The soils analysis evaluates the structure and texture of the soil. The addition of soil amendments can improve the structure and texture of soil, increase its water-holding capacity, and reduce the leaching of fertilizers. Soil amendments, such as compost from clippings, can contribute to an overall healthier plant environment, allowing easier root development and fewer soil-related problems. The use of peat as an amendment should be very limited (such as in containers), as it is both expensive and originates from peat bogs, which are non-renewable. Fertilizers should be applied on the basis of the results of soil tests that have been conducted to identify plant nutritional needs and pH, as described in the "Nutrient Management" chapter.

12.3 Plant Selection

Selection of specific plants should be based to the extent possible on natural ecosystems in the area. This is particularly true for the perimeter zones and out-of-play areas. Native plants can also enhance wildlife habitat by providing forage for different species, such as game and non-game wildlife, birds, and pollinators.

Additional considerations for species selection and placement include design intentions and knowing the ultimate sizes and growth rates of trees, shrubs, and ground covers. This reduces the need for future pruning and debris removal. In addition, the adaptability of plants to a specific site is important. Site-specific characteristics to consider include sun exposure, light intensity, wind conditions, drainage, and temperatures.

The introduction of invasive plants or plants that are potentially invasive should be avoided, and invasive or noxious weed species that are present on the site should be controlled. The SDDA publishes a [noxious weed and pest list](#) and a [weed watch list](#) that can provide useful information for invasive species control efforts.



Figure 37. Coneflowers are a favorite plant of butterflies, bees and other pollinators as well as birds.

12.4 Installation

During landscape bed construction, native soil should be used and any hardpan or compaction from construction should be resolved. The beds should be sloped away from buildings, with a minimum percent slope away from buildings of at least 2% for at least 10 feet. Resolve drainage

issues and establish clear drainage patterns prior to installing plants. Plants with higher moisture requirements can be planted at lower elevations and drought-tolerant plants at higher elevations.

In general, the best times to plant trees and shrubs in South Dakota are in spring or in the early fall. These times reduce the stress on the plants by capitalizing on periods of cooler (but not cold) temperatures and more moisture. However, certain species, including some native plants, perform best if planted only in spring unless a high level of management during establishment is available. Herbaceous plants can also be planted during these time periods, although planting earlier in the fall will give these smaller plants a better chance to establish good root systems.

12.5 Irrigation

Regardless of their ability to tolerate drought, all plants require supplemental irrigation during establishment. To increase water-use efficiency and improve plant establishment in landscaping, consider hand-watering individual plants for the first several months of the growing season. If the plants have been selected and placed in zones that match their water requirements, irrigation can be scheduled to help efficiently meet the water needs of the entire landscape, thus eliminating regular hand-watering. When needed, plants should be watered in the early morning to conserve water by avoiding water loss due to evaporation.

Careful assessment of landscape watering patterns minimizes spray on impervious surfaces, blockage of spray by plants or other obstructions, and runoff on slopes, clay soils, or compacted sites. Focusing the irrigation of woody plants at or beyond the dripline promotes extensive rooting.

For existing irrigation systems, assess the coverage to determine whether changes should be made to identify areas where efficiency can be improved. Ideally, the irrigation system for the landscape beds should be zoned just like the landscape beds. Periodically throughout the growing season, the performance of the landscape irrigation system should be checked.

12.6 Use of Mulch

Mulch conserves soil moisture, mitigates temperature extremes, and reduces weed competition. In winter, mulch helps prevent soil cracks from forming and exposing roots to cold temperatures and winter desiccation. Organic mulches include herbicide-free grass clippings (though avoid applying too deeply to prevent matting and heating the soil) and wood chips of varying dimensions. Organic mulches are preferred, as non-organic mulches such as stone may add heat stress around annuals and perennials.

Annuals and perennials grow best with no more than 2 inches of mulch; mulch around trees and shrubs should be no more than 3-4" deep. With any planting, mulch should be placed between the plants and not on top of the crown or against tree trunks or shrub canes. In winter after the ground freezes, a deeper layer of coarse mulch (evergreen branches) over bulbs and other perennials can delay or prevent early growth.

12.7 Pruning

Correctly pruning trees, shrubs, and herbaceous perennials has multiple benefits throughout a landscape or golf course. Trees and shrubs are pruned first for safety. Pruning in some cases can increase plant health and result in better growth in future seasons. Typically, the ideal time to prune deciduous shade trees in South Dakota is in March to early April, except in times of drought. Shrubs should be pruned based on their season of bloom (if the flowers are significant). Plants that bloom on second-year or old wood set their flower buds immediately after flowering and can be pruned for the month following bloom. Plants that bloom on new wood, or current-season wood, can be pruned in early spring prior to dormancy break. For more information on pruning, see SDDA's [The Basics of Tree Pruning](#).



Figure 38. Tree pruning using a safety harness.

12.8 Pest Management

The same principles and methods identified in the IPM chapter can be applied to landscaped areas. The UNL Extension's [Integrated Pest Management for Landscapes](#) provides guidance specifically for these areas. The SDSU Extension's [2018 Weed Control: Noxious Weeds](#) publication provides information on controlling listed noxious and invasive weeds.



Figure 39. Pine beetle damage.

12.9 Landscape Best Management Practices

Planning and Design

- ❖ Leave the majority of non-play areas in natural vegetation -- the perimeter zone.
- ❖ Enhance natural areas with supplemental plantings of native and adapted species.
- ❖ In landscaped areas, use natural drainage patterns and directional site grading to channel runoff away from impervious surfaces onto planted areas such as grass swales, filter strips, or rain gardens.
- ❖ Install rain gardens in locations where they can catch and temporarily hold runoff.
- ❖ Minimize the amount of area covered by paved surfaces. Where feasible, use permeable materials such as bricks laid on sand, interlocking pavers or pervious pavers, porous concrete, mulch, or plants.
- ❖ Use a zoned approach to plant management and water needs and minimize the areas zoned for high water use.

Site Inventory and Assessment

- ❖ Conduct an inventory of existing plants, their condition and quality, and their contribution to the overall style of the course.
- ❖ Conduct a soil analysis before choosing specific plants for landscape areas.

- ❖ Conduct a soil test before applying fertilizers. Modify pH if needed, based on soil test results.
- ❖ Amend the soil to improve soil texture and increase water infiltration.

Plant Selection

- ❖ Select native species whenever possible; use adapted species or cultivars of native plants where appropriate.
- ❖ Select trees, plants, and grass species to attract birds seeking wild fruits, herbs, seeds, nesting materials, cover, and insects.
- ❖ Know the ultimate sizes and growth rates of trees, shrubs, herbaceous plants, and ground covers.
- ❖ Select plants recommended for your specific location.
- ❖ Choose the most stress-tolerant species for a particular area.
- ❖ Do not introduce invasive species into the landscape.
- ❖ Control or remove existing invasive species and noxious weeds.

Irrigation

- ❖ Irrigate frequently during establishment.
- ❖ Water established plants based on their needs and, when necessary, deeply and infrequently.
- ❖ Irrigate in the early morning to conserve water.
- ❖ Avoid water runoff onto impervious surfaces or slopes.
- ❖ Evaluate landscape irrigation performance periodically.

Use of Mulch

- ❖ Use mulch in landscaped beds.
- ❖ Use organic mulches whenever possible.
- ❖ Use only herbicide-free grass clippings for mulch.
- ❖ Protect bulbs and other perennials in winter with a layer of coarse mulch (evergreen branches) to delay or prevent early growth.

Pruning

- ❖ Hire a certified arborist to prune trees as the correct pruning cuts are essential to good tree health.
- ❖ Maintain pruning equipment to ensure clean cuts and less risk of damage to the plant.
- ❖ Prune deciduous shade trees in March and early April, except in times of extreme drought.
- ❖ Prune shrubs based on their season of bloom.

Pest Management

- ❖ Use IPM for landscaped areas.

13 ENERGY

The use of energy for all activities in society is of great interest worldwide. Golf courses use a variety of energy sources, primarily electricity, gasoline, diesel, natural gas, propane, and heating oil. Renewable sources, such as solar, wind, and geothermal, are increasingly being utilized and considered by small business as the return on investment increases. These newer technologies offer opportunities to reduce dependencies on fossil fuels and our carbon footprint.

To establish effective energy BMPs, the facility's existing energy consumption should be evaluated, and improvements should be achieved through energy reduction, conservation, and new technologies. Energy audits allow for identification of deficiencies. A written energy conservation plan is key to ensuring improvement.

Managers should evaluate current conservation practices based on these categories:

- Buildings, infrastructure, and facility amenities such as the clubhouse, restaurant, kitchen, swimming pool, parking lot, offices, maintenance building(s), tennis courts, restrooms, etc.
- Golf course and surrounding landscapes, and related agronomic operations (playing surfaces, equipment, turfgrass maintenance, etc.).
- Irrigation systems and pump station.

The irrigation system and pump station are the largest consumers of energy on the golf course and should be evaluated. Conserving and reducing water through irrigation BMPs also reduces energy consumption.

Some policies, financial incentives, and loan opportunities exist at the state and local level for renewables and energy efficiency measures at commercial enterprises such as golf courses. Energy providers can provide information, expertise, and incentives to help achieve these goals.

13.1 Energy Audits and Evaluation

An energy audit of the facility should be done if one has not been conducted previously. Energy audits identify areas most in need of conservation. Utility providers can be a source of expertise in conducting an audit. An energy audit should include these steps:

- Evaluate insulation in heated buildings.
- Evaluate heating, ventilation, and air conditioning (HVAC) system efficiency.
- Determine annual energy usage.
- Itemize usage according to various categories.
- Determine if energy usage during non-peak hours are maximized.
- Compare usage with similar small businesses.
- Identify areas of improvement.

Determining energy conservation goals and establishing an environmental plan is a first step in addressing energy efficiency. An energy management plan sets a baseline related to current energy use and incorporates quality management elements (plan, do, check, and act) for continual improvements. Once goals for energy conservation are established and documented, this policy should be communicated to all staff.

Evaluating the performance of an energy conservation program requires tracking and measuring energy use at the facility based on energy assessment units (e.g. kilowatt hour, BTU). Monitoring energy usage can be accomplished with energy management software or programs such as [EPA's Portfolio Manager](#), which also incorporates features such as reporting, savings calculations, and carbon footprint calculations. To benchmark performance, energy consumption can be compared with other local golf facilities of similar size or more generally to buildings of similar size.

13.2 Energy Efficiency Improvements

The audit will identify opportunities to increase energy efficiency in buildings, amenities, and operations. For example, ground-based heat pumps conserve energy as compared with conventional heating sources and could be considered for new building construction or replacement for existing heat sources when the opportunities arise. Developing and implementing a viable energy conservation plan will lead to improvements over time.

13.3 Green and Alternative Energy

Green and alternative energy can be incorporated into golf course operations. Golf courses can become small-scale generators of energy through wind, solar, and photovoltaic installations and geothermal heating and pumping. Golf courses normally have the land, space, and natural resources available on the property to lend themselves to energy generation as newer technologies become more affordable. Financial and tax incentives may be available for installing these energy generators.



Figure 40. Golf courses can become small-scale generators of energy.

13.4 Energy Best Management Practices

Energy Audits and Evaluation

- ❖ Conduct an energy audit, including lighting, insulation, and HVAC systems.
- ❖ Monitor energy use by tracking statistics and “time of use” data.
- ❖ Install precision meters, gauges, etc.
- ❖ Develop an equipment inventory that documents individual equipment’s energy use, traffic patterns, maintenance records, operation hours, etc.
- ❖ Benchmark performance against similar-sized facilities.
- ❖ Educate, train, and motivate employees on energy efficiency practices pertaining to golf course operations.

Energy Efficiency Improvements

- ❖ Evaluate and monitor all energy sources, tracking both costs and any usage trends.
- ❖ Add insulation where needed.
- ❖ Use non-peak electrical hours for charging golf carts and maintenance equipment.
- ❖ Prioritize pump station usage during non-peak hours.
- ❖ Limit high-consumption activities when demand is high.
- ❖ Install LED lighting and other high-efficiency alternatives.
- ❖ Install motion sensors for lights where appropriate.
- ❖ Install low-flow faucets.
- ❖ Install programmable thermostats.
- ❖ Consider energy management software.
- ❖ Utilize the [EPA’s Energy Star and Portfolio Manager](#) programs.
- ❖ Utilize the [EPA’s WaterSense](#) program.
- ❖ Maintain good record-keeping practices.
- ❖ Prioritize energy consumption as part of decision-making when making purchases concerning all aspects of facility management.
- ❖ Evaluate effectiveness of upgrades according to efficiency and conservation goals for energy use.

Green and Alternative Energy

- ❖ Consider pursuing the U.S. Green Building Council’s LEED certification for new buildings and existing building retrofits.
- ❖ Use alternative energy from natural sources, such as solar, geothermal, and wind energy generation when possible.
- ❖ Assess the viability of small-scale wind, solar, and photovoltaic installations.
- ❖ Install geothermal heating and cooling systems, if possible.

REFERENCES

(Note: URLs are current as of July 2019.)

- Arling, Michelle, et al. *National Pesticide Applicator Certification Core Manual, second edition*. National Association of State Departments of Agriculture Research Foundation. 2014.
https://s3.amazonaws.com/nasda2/media/Reports/Core_Title-Page-and-Acknowledgements.pdf?mtime=20171025135711
- Ball, John, and Aaron Kiesz. *The Basics of Tree Pruning*. South Dakota Department of Agriculture.
<https://sdda.sd.gov/legacydocs/Forestry/publications/PDF/The-Basics-of-Tree-Pruning.pdf>
- Bauer, Erin C., Clyde L. Ogg, Jan R. Hygnstrom, Emilee A. Dorn, and Ben W. Beckman. *Pesticide Safety: Choosing the Right Gloves*. University of Nebraska-Lincoln Extension. 2015.
<http://extensionpublications.unl.edu/assets/pdf/g1961.pdf>
- Baum-Haley, Melissa. *2014 Landscape Irrigation Best Management Practices*. Irrigation Association and the American Society of Irrigation Consultants. 2014. Download via:
<https://www.irrigation.org/IA/Advocacy/Standards-Best-Practices/Landscape-Irrigation-BMPs/IA/Advocacy/Landscape-Irrigation-BMPs.aspx>
- Borders, Brianna, et al. *Pollinator Plants of the Central United States Native Milkweeds (Asclepias spp.)*. Xerces Society for Invertebrate Conservation. 2013.
https://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/mopmcpu11905.pdf
- Chalmers, David R. *Buying Quality Grass Seed For Lawns, Parks, and Sports Turf in the Northern Great Plains*. South Dakota State University Extension. 2013.
<https://docplayer.net/29705156-Buying-quality-grass-seed-for-lawns-parks-and-sports-turf-in-the-northern-great-plains.html>
- Dobbs, Emily, and Daniel Potter. "Operation Pollinator for Golf Courses." *Golf Course Management*. April 2013: 100-103.
<http://gcmdigital.gcsaa.org/i/118283-apr-2013/111>
- Duncan, R.R., R.N. Carrow, and M. Huck. *Understanding Water Quality and Guidelines to Management*. United States Golf Association. 2000.
<http://gsrpdf.lib.msu.edu/ticpdf.py?file=/2000s/2000/000914.pdf>
- Emmons, R.D. *Turfgrass Science and Management*, 3rd ed. Delmar Publishers. 2000.
- EnviroLogic Resources. *OGCSA Environmental Stewardship Guidelines, Second Edition*. Oregon Golf Course Superintendents' Association. 2009.
<https://www.gcsaa.org/uploadedfiles/Environment/Get-Started/BMPs/Oregon-Chapter-Environmental-Stewardship-Guidelines.pdf>
- Frogge, Mary Jane, and Soni Cochran. *Attracting Pollinators to Your Landscape*. University of Nebraska-Lincoln Extension.
<https://lancaster.unl.edu/pest/resources/354Pollinators.pdf>
- Giesler, Loren J. *Dollar Spot Disease of Turfgrass*. University of Nebraska-Lincoln Extension. 2009.
<https://turf.unl.edu/NebGuides/dollarspot.pdf>
- Giesler, Loren J. *Pythium Blight of Turfgrass*. University of Nebraska-Lincoln Extension. 2009.
<https://turf.unl.edu/NebGuides/Pythium.pdf>
- Hygnstrom, Jan R., Clyde L. Ogg, and Cheryl A. Alberts. *Respirators for Handling Pesticides*. University of Nebraska-Lincoln Extension. 2017.
<http://extensionpublications.unl.edu/assets/pdf/ec3021.pdf>

- Hygnstrom, Jan R., Clyde L. Ogg, Erin C. Bauer, Pierce J. Hansen, and Anne M. Streich. *Integrated Pest Management for Landscapes*. University of Nebraska-Lincoln Extension. 2013.
<http://extensionpublications.unl.edu/assets/pdf/ec1266.pdf>
- Iles, Derric L. "South Dakota's Aquifers." *Quality on Tap!* October 2008: 8-9.
http://www.sdgs.usd.edu/pdf/SD_Aquifers_article.pdf
- Johnson, Bill, Bill Casaday, Dallas Peterson, and Dennis Kuhlman. *Cleaning Fields Sprayers to Avoid Crop Injury*. University of Missouri-Columbia Extension. 1997.
<https://extensiondata.missouri.edu/pub/pdf/agguides/crops/g04852.pdf>
- Johnson, Paul O., David Vos, Jill Alms, and Leon Wrage. *2018 Weed Control: Noxious Weeds*. 2018.
[https://www.sdgs.usd.edu/pdf/2018_Weed_Control_Noxious_Weeds](https://www.sdgs.usd.edu/pdf/2018_Weed_Control_Noxious_Weeds.pdf)
- Johnson, Rex R., Kenneth F. Higgins, Michael L. Kjellsen, and Charles R. Elliott. *Eastern South Dakota Wetlands*. South Dakota State University. 1997.
<https://www.fws.gov/wetlands/Documents/Eastern-South-Dakota-Wetlands.pdf>
- Kreuser, Bill C. *Simplifying Soil Test Interpretations for Turf Professionals*. University of Nebraska-Lincoln Extension. 2015.
<http://extensionpublications.unl.edu/assets/pdf/g2265.pdf>
- Kruger, Greg R., Robert N. Klein, and Clyde L. Ogg. *Spray Drift of Pesticides*. University of Nebraska-Lincoln Extension. 2013.
<http://extensionpublications.unl.edu/assets/pdf/g1773.pdf>
- Landschoot, Peter. *Irrigation Water Quality Guidelines for Turfgrass Sites*. Penn State University.
<http://plantscience.psu.edu/research/centers/turf/extension/factsheets/water-quality>
- Larson, Jonathan, David Held, and R. Chris Williamson. *Best Management Practices for Turf Care and Pollinator Conservation: Fast Facts*. USDA National Institute of Food and Agriculture. 2016.
<https://www.ncipmc.org/projects/publications/best-management-practices-for-turf-care-and-pollinator-conservation-fast-facts/>
- Ley, Elizabeth L. *Selecting Plants for Pollinators: A Regional Guide for Farmers, Land Managers, and Gardeners in the Prairie Parkland*. North American Pollinator Protection Campaign and Pollinator Partnership.
<https://www.pollinator.org/PDFs/PrairieParkland.rx11.pdf>
- Moisset, Beatriz, and Stephen Buchmann. *Bee Basics: An Introduction to Our Native Bees*. USDA Forest Service and Pollinator Partnership. 2011.
https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5306468.pdf
- Nebraska Game and Parks Commission. *Resolving Common Maintenance Problems/Aquatic Vegetation*. Nebraska Game and Parks Commission. 2015.
https://outdoornebraska.gov/wp-content/uploads/2015/11/PMGS_Aquatic_Vegetation.pdf
- Ogg, Barb. *Managing Canada Geese*. University of Nebraska-Lincoln Extension. 2009.
<https://lancaster.unl.edu/pest/resources/337cangeese.pdf>
- Ogg, Clyde L., Jan R. Hygnstrom, Cheryl A. Alberts, and Erin C. Bauer. *Managing the Risk of Pesticide Poisoning and Understanding the Signs and Symptoms*. University of Nebraska-Lincoln Extension. 2018.
<http://extensionpublications.unl.edu/assets/pdf/ec2505.pdf>
- Ogg, Clyde L., et al. *Protective Clothing and Equipment for Pesticide Applicators*. University of Nebraska-Lincoln Extension. 2018.
<http://extensionpublications.unl.edu/assets/pdf/g758.pdf>
- Puckett, Greg J., Clyde L. Ogg, Robert N. Klein, and Cheryl A. Alberts. *Cleaning Pesticide Application Equipment*. University of Nebraska-Lincoln Extension. 2018.
<http://extensionpublications.unl.edu/assets/pdf/g1770.pdf>

[Purdue University Extension. *Turfgrass Weed Control for Professionals, 2017 edition*. Purdue University Extension. 2017.](#)

<https://marketplace.unl.edu/extension/turfgrass-weed-control-for-professionals.html>

Sallenave, Rossana. [Stream Biomonitoring Using Macrobenthic Invertebrates](#)

New Mexico State University Cooperative Extension Service. 2015.

http://aces.nmsu.edu/pubs/_circulars/CR677.pdf

Schild, Jim A., and Scott Dworak. [Water Wise: Drought Effects on Turf in the Landscape](#). University of Nebraska-Lincoln Extension. 2013.

<http://extensionpublications.unl.edu/assets/pdf/g2191.pdf>

Shepherd, Matthew. [Making Room for Native Pollinators How to Create Habitat for Pollinator Insects on Golf Courses](#). United States Golf Association. 2002.

http://www.xerces.org/wp-content/uploads/2008/10/making_room_for_pollinators_usga1.pdf

South Dakota Department of Agriculture. [Fertilizer and Pesticide Bulk Facility Manual](#). South Dakota Department of Agriculture. 2015.

https://sdda.sd.gov/ag-services/pesticide-program/Pesticide%20PDFs_Forms/bulkman_finalupdated.pdf

South Dakota Department of Agriculture. [Managed Pollinator Plan](#). South Dakota Department of Agriculture. 2017.

<https://sdda.sd.gov/ag-services/beekeeping-apiary-resources/pdf/Pollinator.Plan.July2017.pdf>

South Dakota Department of Agriculture. [On-Farm Fertilizer Storage](#). South Dakota Department of Agriculture.

https://sdda.sd.gov/legacydocs/Ag_Services/Agronomy_Services_Programs/Fertilizer_Soil_Amendment_Program/onfarmfertstorage.pdf

South Dakota Department of Agriculture. [Private Applicator Handling and Discharge Response Plan](#). South Dakota Department of Agriculture.

https://sdda.sd.gov/legacydocs/Ag_Services/forms/Priv_PHDRP.pdf

South Dakota Department of Agriculture. [Reducing Pesticide Waste in South Dakota](#). South Dakota Department of Agriculture. 2010.

<https://sdda.sd.gov/documents/farming-ranching-agribusiness/container-recycling-waste-pesticide/reducingpesticidewasteinsouthdakota.pdf>

South Dakota Department of Agriculture. [Riparian Area Management](#). South Dakota Department of Agriculture. 2007.

<https://sdda.sd.gov/legacydocs/Forestry/publications/PDF/Riparian-Area-Management.pdf>

South Dakota Department of Environment and Natural Resources. [Petroleum Assessment and Cleanup Handbook](#).

South Dakota Department of Environment and Natural Resources. 2003.

https://denr.sd.gov/des/gw/Spills/Handbook/Hand_Book.aspx

Stange, Craig. [Rain Gardens: Capturing and Using the Rains of the Great Plains](#). USDA Natural Resources Conservation Service. 2007.

https://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/ndpmctn7278.pdf

Streich, Anne M., Roch E. Gaussoin, and Zac J. Reicher. [Managing Turf and Landscape Weeds](#). University of Nebraska-Lincoln Extension. 2014.

<http://extensionpublications.unl.edu/assets/pdf/ec1269.pdf>

U.S. Fish & Wildlife Service's South Dakota Ecological Services. [South Dakota Listed Species by County List](#). U.S. Fish & Wildlife Service. 2016.

https://www.fws.gov/southdakotafieldoffice/SpeciesByCounty_April2016.pdf

United States Golf Association. [*A Guide to Constructing the USGA Putting Green*](#). United States Golf Association. 2015.

<https://cloud.3dissue.com/73035/73358/87210/CreatingTheUSGAPuttingGreen/html5/index.html>

Vaughan, Mace, Eric Lee-Mader, Sarah Foltz Jordan, Emily Krafft, and Sara Morris. [*Pollinator Meadow Upper Midwest Installation Guide and Checklist*](#). Xerces Society for Invertebrate Conservation. 2015.

http://www.xerces.org/wp-content/uploads/2013/01/InstallGuideJobSheet_UpperMidwest_CnsrvCvr.pdf

Virginia Golf Course Superintendents Association. [*Environmental Best Management Practices for Virginia's Golf Courses*](#). Virginia Cooperative Extension. 2012.

https://pubs.ext.vt.edu/ANR/ANR-48/ANR-48_pdf.pdf

Watkins, E., et al. "[*Low-Input Turfgrass Species for the North Central United States*](#)." January 2011. *Applied Turfgrass Science* 8. doi:10.1094/ATS-2011-0126-02-RS.

<https://mgcsa.org/resources/Documents/Environmental%202016/Low%20input%20turfgrass%20species%20for%20NC%20US.pdf>

Wilson, Jim. [*Waste Pesticides: Proper Storage, Handling, and Disposal*](#). South Dakota State University Cooperative Extension Service. 2002.

https://sdda.sd.gov/documents/farming-ranching-agribusiness/container-recycling-waste-pesticide/SDSU_waste_pesticides.pdf

BIBLIOGRAPHY

Notes: URLs are as of September 2016

- Aerts, M.O., N. Nesheim, and F. M. Fishel. April 1998; revised September 2015. *Pesticide recordkeeping*. PI-20. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI012>.
- Aquatic Ecosystem Restoration Foundation. 2014. *Biology and Control of Aquatic Plants: A Best Management Practices Handbook: 3rd Ed.* Gettys, L.A., W. T. Haller, and D. G. Petty, editors. <http://www.aquatics.org/bmp%203rd%20edition.pdf>
- ASCE, January 2005. *The ASCE standardized reference evapotranspiration equation*. Final report of the Task Committee on Standardization of Reference Evapotranspiration, Environmental and Water Resources Institute of the American Society of Civil Engineers. 1801 Alexander Bell Drive, Reston, VA 20191 Available: <http://www.kimberly.uidaho.edu/water/asceewri/ascestzdetmain2005.pdf>
- Bohmont, B. 1981. *The new pesticide users guide*. Fort Collins, Colorado: B & K Enterprises.
- Brecke, B.J., and J.B. Unruh. May 1991; revised February 25, 2003. *Spray additives and pesticide formulations*. Fact Sheet ENH-82. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/LH061>.
- Broder, M.F., and D.T. Nguyen. 1995. *Coating concrete secondary containment structures exposed to agrichemicals*. Circular Z-361. Muscle Shoals, Alabama: Tennessee Valley Authority, Environmental Research Center. Tel. (205) 386-2714.
- Broder, M.F., and T. Samples. 2002. *Tennessee handbook for golf course environmental management*. Tennessee Department of Agriculture.
- Buss, E.A. January 2002; revised July 2003. *Insect pest management on golf courses*. ENY-351. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/IN410>.
- Butler, T., W. Martinkovic, and O.N. Nesheim. June 1993; revised April 1998. *Factors influencing pesticide movement to groundwater*. PI2. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI002>.
- California Fertilizer Association. 1985. *Western fertilizer handbook*, 7th ed. Sacramento, California.
- Carrow, R.N., R. Duncan, and C. Waltz. 2007. *Best Management Practices (BMPs) Water-Use Efficiency/Conservation Plan for Golf Courses*. Available: [https://www.gcsaa.org/uploadedfiles/Environment/Get-Started/BMPs/Water-use-efficiency-and-conservation-best-management-practices-\(Georgia\).pdf](https://www.gcsaa.org/uploadedfiles/Environment/Get-Started/BMPs/Water-use-efficiency-and-conservation-best-management-practices-(Georgia).pdf)
- Carrow, R.N., R.R. Duncan, and D. Wienecke. 2005. BMPs: Critical for the golf industry. *Golf Course Management*. 73(6):81-84.
- Center for Resource Management. 1996. *Environmental principles for golf courses in the United States*. 1104 East Ashton Avenue, Suite 210, Salt Lake City, Utah 84106. Tel: (801) 466-3600, Fax: (801) 466-3600.
- Clark, G.A. July 1994. *Microirrigation in the landscape*. Fact Sheet AE254. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/AE076>.
- Clark, Mark and Acomb, Glenn; Florida Field Guide to Low Impact Development: Stormwater Reuse. Univ. Florida 2008. http://buildgreen.ufl.edu/Fact_sheet_Stormwater_Reuse.pdf
- Colorado Nonpoint Source Task Force. 1996. *Guidelines for Water Quality Enhancement at Golf Courses Through the Use of Best Management Practices*. Available: <http://www.wrightwater.com/assets/7-golf-course-bmps.pdf>

Connecticut Department of Environmental Protection. 2006. Best Management Practices for Golf Course Water Use. Available: http://www.ct.gov/deep/lib/deep/water_inland/diversions/golfcoursewaterusebmp.pdf

Cromwell, R.P. June 1993; reviewed December 2005. *Agricultural chemical drift and its control*. CIR1105. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/AE043>.

Crow, W.T. February 2001; revised November 2005. *Nematode management for golf courses in Florida*. ENY-008 (IN124). Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/IN124>.

Daum, D.R., and T.F. Reed. n.d. *Sprayer nozzles*. Ithaca, New York: Cornell Cooperative Extension. Available <http://psep.cce.cornell.edu/facts-slides-self/facts/gen-peapp-spray-nozz.aspx>.

Dean, T.W. February 2003. *Pesticide applicator update: Choosing suitable personal protective equipment*. PI-28. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI061>.

———. April 2004; revised November 2004. *Secure pesticide storage: Facility size and location*. Fact Sheet PI-29. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI064>.

———. April 2004; revised November 2004. *Secure pesticide storage: Essential structural features of a storage building*. Fact Sheet PI-30. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI065>.

Dean, T.W., O.N. Nesheim, and F. Fishel. Revised May 2005. *Pesticide container rinsing*. PI-3. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI003>.

Delaware Nutrient Management Commission. 2006. Water Quality Best Management Practices: Nutrients, Irrigation and Pesticides for Golf Course, Athletic Turf, Lawn Care and Landscape Industries. Available: <http://dda.delaware.gov/nutrients/forms/BMPnonagforprinter.pdf>

Dodson, R.G. 2000. *Managing wildlife habitat on golf courses*. Sleeping Bear Press. Chelsea, MI.

Elliott, M.L., and G.W. Simone. July 1991; revised April 2001. *Turfgrass disease management*. SS-PLP-14. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/LH040>.

Fishel, F.M. March 2005. *Interpreting pesticide label wording*. Gainesville, Florida: Institute of Food and Agricultural Sciences. Available: <http://edis.ifas.ufl.edu/PI071>.

Fishel, F.M., and Nesheim, O.N. November 2006. *Pesticide safety*. FS11. Gainesville, Florida: Institute of Food and Agricultural Sciences. Available: <http://edis.ifas.ufl.edu/pdf/CV/CV10800.pdf>.

Florida Department of Agriculture and Consumer Services. n.d. *Pesticide recordkeeping—benefits and requirements*. Available: <http://www.flaes.org/pdf/Pesticide%20Recordkeeping%20Pamphlet%205-05.pdf>.

Florida Department of Agriculture and Consumer Services. Division of Agricultural Environmental Services. *Suggested pesticide recordkeeping form*. Available: <https://www.freshfromflorida.com/content/download/2990/18861/Suggested%20Pesticide%20Recordkeeping%20Form.pdf>

———. Division of Agricultural Environmental Services. *Suggested pesticide recordkeeping form for organo-auxin herbicides*. Available: <http://forms.freshfromflorida.com/13328.pdf>.

Florida Department of Agriculture and Consumer Services and Florida Department of Environmental Protection. 1998. *Best management practices for agrichemical handling and farm equipment maintenance*. Available: <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/agbmp3p.pdf>

Florida Department of Environmental Protection. 2008. *Florida stormwater, erosion, and sedimentation control inspector's manual*. Tallahassee, Florida: Nonpoint Source Management Section, MS 3570, 3900 Commonwealth Blvd., Tallahassee, Florida 32399-3000. Available: <http://www.dep.state.fl.us/water/nonpoint/docs/erosion/erosion-inspectors-manual.pdf>.

———. December 27, 2002. *Environmental risks from use of organic arsenical herbicides at south Florida golf courses*. FDEP white paper. Available: <http://fdep.ifas.ufl.edu/msma.htm>.

———. April 2002. *Florida water conservation initiative*. Available: http://www.dep.state.fl.us/water/waterpolicy/docs/WCI_2002_Final_Report.pdf.

———. 2015. "Florida-friendly Best Management Practices for Protection of Water Resources by the Green Industries", Florida Department of Environmental Protection. Revised December 2008, 3rd printing 2015. <https://fyn.ifas.ufl.edu/pdf/grn-ind-bmp-en-12-2008.pdf>

———. 2012. *Best Management Practices for The Enhancement of Environmental Quality on Florida Golf Courses*. Florida Department of Environmental Protection. 3rd printing, September 2012. <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glbmp07.pdf>

———. Revised August 2009. *A guide on hazardous waste management for Florida's auto repair shops*. Available: http://www.dep.state.fl.us/waste/quick_topics/publications/shw/hazardous/business/Paint_and_Body8_09.pdf.

———. October 2005. *Checklist guide for 100% closed loop recycle systems at vehicle and other equipment wash facilities*. Available: <http://www.dep.state.fl.us/water/wastewater/docs/ChecklistGuideClosed-LoopRecycleSystems.pdf>.

———. October 2005. *Guide to best management practices for 100% closed-loop recycle systems at vehicle and other equipment wash facilities*. Pollution Prevention Program and Industrial Wastewater Section. Available: <http://www.dep.state.fl.us/water/wastewater/docs/GuideBMPClosed-LoopRecycleSystems.pdf>.

———. 2006. *State of Florida erosion and sediment control designer and reviewer manual*. Nonpoint Source Management Section. Available: <http://www.dep.state.fl.us/water/nonpoint/erosion.htm>.

———. 2016. *Operation Cleansweep for Pesticides Web site*. Available: <http://www.dep.state.fl.us/waste/categories/cleansweep-pesticides>.

———. December 1, 2005. *Standards and specifications for turf and landscape irrigation systems*, 5th Ed. Available: <http://ufdc.ufl.edu/UF00076845/00001>.

———. December 2006. *Landscape Irrigation & Florida-Friendly Design Standards*. Florida Department of Environmental Protection, Office of Water Policy, 3900 Commonwealth Blvd., MS 46, Tallahassee, FL 32399-3000. Available: <http://www.dep.state.fl.us/water/waterpolicy/docs/LandscapeIrrigationFloridaFriendlyDesign.pdf>

Gilman, E. 2006. *Pruning shade trees in landscapes*. Available: <http://hort.ufl.edu/woody/pruning/index.htm>.

Golf Course Superintendents Association of America. 2012. *Golf Course Environmental Profile; Volume IV; Energy Use and Energy Conservation Practices on U.S. Golf Courses*. Available: <https://www.gcsaa.org/Uploadedfiles/Environment/Environmental-Profile/Energy/Golf-Course-Environmental-Profile--Energy-Use-and-Conservation-Report.pdf>

Golf Course Water Resources Handbook of Best Management Practices (Pennsylvania). 2009. Available: <http://pecpa.org/wp-content/uploads/Golf-Course-Water-Resources-Handbook-of-Best-Management-Practices.pdf>

Havlin, J.L., et al. 2004. *Soil fertility and fertilizers*, 7th Ed. Prentice Hall.

Haydu, J.J., and A.W. Hodges. 2002. *Economic impacts of the Florida golf course industry*. UF-IFAS Report EIR 02-4. Available: <http://economicimpact.ifas.ufl.edu/publications/EIR02-4r.pdf>.

Helfrich, L.A., et al. June 1996. *Pesticides and aquatic animals: A guide to reducing impacts on aquatic systems*. Virginia Cooperative Extension Service. Publication Number 420-013. Available: <http://www.ext.vt.edu/pubs/waterquality/420-013/420-013.html>.

- Hornsby, A.G., T.M. Buttler, L.B. McCarty, D.E. Short, R.A. Dunn, G.W. Simone. Revised September 1995. *Managing pesticides for sod production and water quality protection*. Circular 1012. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/SS053>.
- Insecticide Resistance Action Committee Web site. Available: <http://www.irac-online.org/>.
- King, K.W., and J.C. Balogh. 2001. Water quality impacts associated with converting farmland and forests to turfgrass. In: *Transactions of the ASAE, Vol. 44(3): 569-576*.
- Lehtola, C.J., C.M. Brown, and W.J. Becker. November 2001. *Personal protective equipment. OSHA Standards 1910.132-137*. AE271. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/OA034>.
- McCarty, L.B., and D.L. Colvin. 1990. *Weeds of southern turfgrasses*. Gainesville, Florida: University of Florida.
- Midwest Plan Service. Revised 1995. *Designing facilities for pesticide and fertilizer containment*. MWPS-37. Midwest Plan Service, 122 Davidson Hall, Iowa State University, Ames, IA 50011-3080. Tel.: (515) 294-4337. Available: <http://infohouse.p2ric.org/ref/50/49471.pdf>.
- Mitra, S. 2006. *Effects of recycled water on turfgrass quality maintained under golf course fairway conditions*. WaterReuse Foundation, 1199 North Fairfax Street, Suite 410, Alexandria, VA 22314. Available: <http://www.watereuse.org/Foundation/documents/wrf-04-002.pdf>.
- National Pesticide Telecommunications Network. December 1999. *Signal words*. Fact Sheet. Available: <http://npic.orst.edu/factsheets/signalwords.pdf>.
- Nesheim, O.N., and F.M. Fishel September 2007, reviewed August 2013. *Interpreting PPE statements on pesticide labels*. P116. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <https://edis.ifas.ufl.edu/pdf/PI/CV/CV28500.pdf>.
- Nesheim, O.N., and F.M. Fishel. March 1989; revised November 2005. *Proper disposal of pesticide waste*. PI-18. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/PI010>.
- Nesheim, O.N., F.M. Fishel, and M. Mossler. July 1993. *Toxicity of pesticides*. PI-13. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/pdf/PI/PI00800.pdf>.
- O'Brien, P. July/August 1996. Optimizing the turfgrass canopy environment with fans. *USGA Green Section Record, Vol. 34(4), 9-12* Available: <http://gsrpdf.lib.msu.edu/ticpdf.py?file=/1990s/1996/960709.pdf>.
- O'Brien, P., and C. Hartwiger. March/April 2003. Aerification and sand topdressing for the 21st century. *USGA Green Section Record, Vol. 41(2), 1-7*. Available: <http://turf.lib.msu.edu/2000s/2003/030301.pdf>.
- Olexa, M.T., A. Leviten, and K. Samek. December 2008, revised December 2013. *Florida solid and hazardous waste regulation handbook: Table of contents*. FE758. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/fe758>.
- Otterbine Barebo, Inc. 2003. *Pond and lake management*. 3840 Main Road East, Emmaus, PA 18049. Available: <http://www.otterbine.com/assets/base/resources/PondAndLakeManual.pdf>.
- Peterson, A. 2000. *Protocols for an IPM system on golf courses*. University of Massachusetts Extension Turf Program.
- Pennsylvania Department of Environmental Protection, LandStudies, Inc., The Pennsylvania Environmental Council. *Golf Course Water Resources Handbook of Best Management Practices*. June 2009. <http://pecpa.org/wp-content/uploads/Golf-Course-Water-Resources-Handbook-of-Best-Management-Practices.pdf>
- Pettinger, N.A. 1935. Useful chart for teaching the relation of soil reaction to availability of plant nutrients to crops. *Virginia Agri. Ext. Bul. 136, 1-19*.

- Portness, R.E., J.A. Grant, B. Jordan, A.M. Petrovic, and F.S. Rossi. 2014. Best Management Practices for New York State Golf Courses. Cornell Univ. Available: http://nysgolfbmp.cals.cornell.edu/ny_bmp_feb2014.pdf
- Rao, P.S.C., and A.G. Hornsby. May 1993; revised December 2001. *Behavior of pesticides in soils and water*. Fact Sheet SL40. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/SS111>.
- Rao, P.S.C., R.S. Mansell, L.B. Baldwin, and M.F. Laurent. n.d. *Pesticides and their behavior in soil and water*. Ithaca, New York: Cornell Cooperative Extension. Available: <http://psep.cce.cornell.edu/facts-slides-self/facts/gen-pubre-soil-water.aspx>.
- Rodgers, J. n.d. *Plants for lakefront revegetation*. Invasive Plant Management, Florida Department of Environmental Protection, 3900 Commonwealth Blvd., MS 705, Tallahassee, FL 32399. Available: <http://myfwc.com/media/2518526/LakefrontRevegetation.pdf>.
- Sartain, J.B. 2000. *General recommendations for fertilization of turfgrasses on Florida soils*. Fact Sheet SL-21. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/LH014>.
- . 2001. *Soil testing and interpretation for Florida turfgrasses*. SL-181. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/SS317>.
- . 2002. revised October 2006. *Recommendations for N, P, K, and Mg for golf course and athletic field fertilization based on Mehlich-I extractant*. SL-191. Available: <http://edis.ifas.ufl.edu/SS404>. Gainesville, Florida.
- Sartain, J.B., and W.R. Cox. 1998. *The Florida fertilizer label*. SL-3. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/SS170>.
- Sartain, J.B., G.L. Miller, G.H. Snyder, and J.L. Cisar. 1999a. Plant nutrition and turf fertilizers. In: J.B. Unruh and M. Elliott (Eds.). *Best management practices for Florida golf courses*. SP-141 2nd ed. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida.
- . 1999b. Liquid fertilization and foliar feeding. In: J.B. Unruh and M. Elliott (Eds.), *Best management practices for Florida golf courses*. SP-141 2nd ed. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida.
- Sartain, J.B., G.L. Miller, G.H. Snyder, J.L. Cisar, and J.B. Unruh. 1999. Fertilization programs. In: J.B. Unruh and M. Elliott (Eds.). *Best management practices for Florida golf courses*. SP-141 2nd ed. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida.
- Schueler, T.R. 2000. Minimizing the impact of golf courses on streams. Article 134 in: *The practice of watershed protection*. T. R. Schueler and H. K. Holland (Eds.). Ellicott City, Maryland: Center for Watershed Protection. Available: <http://www.stormwatercenter.net/>.
- Schumann, G.L., et al. January 1998. *IPM handbook for golf courses*. Indianapolis, Indiana: Wiley Publishing, Inc.
- Seelig, B. July 1996. *Improved pesticide application BMP for groundwater protection from pesticides*. AE-1113. Fargo, North Dakota: North Dakota State University Extension Service. Available: <http://www.ext.nodak.edu/extpubs/h2oqual/watgrnd/ae1113w.htm>.
- Smajstrla, A.G., and B.J. Boman. April 2000. *Flushing procedures for microirrigation systems*. Bulletin 333. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/WI013>.
- Staples, A.J. 2. Golf Course Energy Use Part 2: Pump Stations. Golf Course Management, July 2009. <https://www.gcsaa.org/Uploadedfiles/Environment/Resources/Energy-Conservation/Golf-course-energy-use-Part-2-Pump-stations.pdf>
- Tennessee Department of Agriculture. Tennessee Handbook for Golf Course Environmental Management. Available: <http://tennesseeturf.utk.edu/pdf/files/golfcourseenvironmentgmt.pdf>

- Thostenson, A., C. Ogg, K. Schaefer, M. Wiesbrook, J. Stone, and D. Herzfeld. 2016. Laundering pesticide-contaminated work clothes. PS 1778. Fargo, ND. North Dakota State Univ. Cooperative Extension. <https://www.ag.ndsu.edu/pubs/plantsci/pests/ps1778.pdf>
- Trautmann, N.M., K.S. Porter, and R.J. Wagenet. n.d. *Pesticides and groundwater: A guide for the pesticide user*. Fact Sheet. Ithaca, New York: Cornell Cooperative Extension. Available: <http://psep.cce.cornell.edu/facts-slides-self/facts/pest-gr-gud-grw89.aspx>
- University of Florida—Institute of Food and Agricultural Sciences. Center for Aquatic and Invasive Plants Web site. Available: <http://plants.ifas.ufl.edu/>.
- . Insect Identification Service Web site. Available: <http://edis.ifas.ufl.edu/SR010>.
- . Nematode Assay Laboratory Web site. Available: <http://edis.ifas.ufl.edu/SR011>.
- . Pesticide Information Office Web site. Available: <http://pested.ifas.ufl.edu/>
- . Plant Disease Clinic Web site. Available: <http://plantpath.ifas.ufl.edu/extension/plant-diagnostic-center/>
- . Rapid Turfgrass Diagnostic Service Web site. Available: <http://turfpath.ifas.ufl.edu/rapiddiag.shtml>.
- Unruh, J.B. November 1993. *Pesticide calibration formulas and information*. Fact Sheet ENH-90. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://edis.ifas.ufl.edu/WG067>.
- Unruh, J.B. 2006. *2006 University of Florida's pest control guide for turfgrass managers*. Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida. Available: <http://turf.ufl.edu>.
- Unruh, J.B., and B.J. Brecke. Revised January 1998. *Response of turfgrass and turfgrass weeds to herbicides*. ENH-100. Gainesville, Florida: Department of Environmental Horticulture, University of Florida. Available: <http://edis.ifas.ufl.edu/WG071>.
- Unruh, J.B., and M. Elliot. 1999. *Best management practices for Florida golf courses*, 2nd ed. UF–IFAS Publication SP-141. Gainesville, Florida.
- Unruh, J.B., J.L. Cisar, and G.L. Miller. 1999. Mowing. In: J.B. Unruh and M.L. Elliot (Eds.). *Best management practices for Florida golf courses*, 2nd ed. Gainesville, Florida: University of Florida Institute of Food and Agricultural Sciences.
- Unruh, J.B., A.E. Dudeck, J.L. Cisar, and G.L. Miller. 1999. Turfgrass cultivation practices. In: J.B. Unruh and M.L. Elliot (Eds.). *Best management practices for Florida golf courses*, 2nd ed. Gainesville, Florida: University of Florida Institute of Food and Agricultural Sciences.
- U.S. Environmental Protection Agency. 2005. *GreenScapes: Environmentally beneficial landscaping*; Washington, D.C. Office of Solid Waste and Emergency Response. Available: <https://archive.epa.gov/greenbuilding/web/pdf/brochure.pdf>
- United States Golf Association. 2004. *Recommendations for a method of putting green construction*. Available: <http://www.usga.org/content/dam/usga/images/course-care/2004%20USGA%20Recommendations%20For%20a%20Method%20of%20Putting%20Green%20Cons.pdf>.
- van Es., H.M. October 1990. *Pesticide management for water quality: Principles and practices*. October 1990. Ithaca, New York: Cornell Cooperative Extension. Available: <http://psep.cce.cornell.edu/facts-slides-self/facts/pestmgmt-water-qual-90.aspx>.
- Virginia Golf Course Superintendents Association. 2012. Environmental Best Management Practices for Virginia's Golf Courses. https://pubs.ext.vt.edu/ANR/ANR-48/ANR-48_pdf.pdf
- White, C.B. 2000. *Turfgrass manager's handbook for golf course construction, renovation, and grow-in*. Sleeping Bear Press. Chelsea, MI.

Best Management Practices for South Dakota Golf Courses

Witt, J.M. n.d. *Agricultural spray adjuvants*. Ithaca, New York: Cornell Cooperative Extension. Available: <http://pmep.cce.cornell.edu/facts-slides-self/facts/gen-peapp-adjuvants.html>.

Yergert, M.B. Austin, and R. Waskom. June 1993. *Best management practices for turfgrass production*. Turf BMP Fact Sheet. Colorado Department of Agriculture. Agricultural Chemicals and Groundwater Protection Program. Available: http://hermes.cde.state.co.us/drupal/islandora/object/co%3A3063/datastream/OBJ/download/Best_management_practices_for_turfgrass_production.pdf.